

# Highly entangled quantum spin chains and their extensions by semigroups

Fumihiko Sugino

Center for Theoretical Physics of the Universe, Institute for Basic Science

Workshop on “Matrix Models for Noncommutative Geometry  
and String Theory”

Erwin Schrödinger Institute (ESI), July 12, 2018

# Highly entangled quantum spin chains and their extensions by semigroups

Fumihiko Sugino

Center for Theoretical Physics of the Universe, Institute for Basic Science

Workshop on “Matrix Models for Noncommutative Geometry  
and String Theory”

Erwin Schrödinger Institute (ESI), July 12, 2018

Bravyi et al, Phys. Rev. Lett. **118** (2012) 207202, arXiv: 1203.5801

R. Movassagh and P. Shor, Proc. Natl. Acad. Sci. **113** (2016) 13278,  
arXiv: 1408.1657

F.S. and P. Padmanabhan, J. Stat. Mech. **1801** (2018) 013101,  
arXiv: 1710.10426

P. Padmanabhan, F.S. and V. Korepin, arXiv: 1804.00978

F.S. and V. Korepin, arXiv:1806.04049

# Outline

Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

Rényi entropy

Rényi entropy of Motzkin model

Summary and discussion

# Introduction 1

## Quantum entanglement

- ▶ Most surprising feature of quantum mechanics,  
No analog in classical mechanics

# Introduction 1

## Quantum entanglement

- ▶ Most surprising feature of quantum mechanics, No analog in classical mechanics
- ▶ From pure state of the full system  $S$ :  $\rho = |\psi\rangle\langle\psi|$ , reduced density matrix of a subsystem  $A$ :  $\rho_A = \text{Tr}_{S-A} \rho$  can become mixed states, and has nonzero entanglement entropy

$$S_A = -\text{Tr}_A [\rho_A \ln \rho_A].$$

This is purely a quantum property.

# Introduction 2

## Area law of entanglement entropy

- ▶ Ground states of quantum many-body systems **with local interactions** typically exhibit the area law behavior of the entanglement entropy:  $S_A \propto (\text{area of } A)$
- ▶ Gapped systems in 1D are proven to obey the area law.  
[Hastings 2007]

# Introduction 2

## Area law of entanglement entropy

- ▶ Ground states of quantum many-body systems **with local interactions** typically exhibit the area law behavior of the entanglement entropy:  $S_A \propto (\text{area of } A)$
- ▶ Gapped systems in 1D are proven to obey the area law.  
[Hastings 2007] (Area law violation)  $\Rightarrow$  Gapless
- ▶ For gapless case, (1 + 1)-dimensional CFT violates logarithmically:  $S_A = \frac{c}{3} \ln(\text{volume of } A)$ . [Calabrese, Cardy 2009]

# Introduction 2

## Area law of entanglement entropy

- ▶ Ground states of quantum many-body systems **with local interactions** typically exhibit the area law behavior of the entanglement entropy:  $S_A \propto (\text{area of } A)$
- ▶ Gapped systems in 1D are proven to obey the area law.  
[Hastings 2007] (Area law violation)  $\Rightarrow$  Gapless
- ▶ For gapless case,  $(1 + 1)$ -dimensional CFT violates logarithmically:  $S_A = \frac{c}{3} \ln(\text{volume of } A)$ . [Calabrese, Cardy 2009]
- ▶ Belief for gapless case in  $D$ -dim. (over two decades) :  
 $S_A = O(L^{D-1} \ln L)$  ( $L$ : length scale of  $A$ )



# Introduction 2

## Area law of entanglement entropy

- ▶ Ground states of quantum many-body systems **with local interactions** typically exhibit the area law behavior of the entanglement entropy:  $S_A \propto (\text{area of } A)$
  - ▶ Gapped systems in 1D are proven to obey the area law.  
[Hastings 2007] (Area law violation)  $\Rightarrow$  Gapless
  - ▶ For gapless case, (1 + 1)-dimensional CFT violates logarithmically:  $S_A = \frac{c}{3} \ln(\text{volume of } A)$ . [Calabrese, Cardy 2009]
  - ▶ Belief for gapless case in  $D$ -dim. (over two decades) :  
 $S_A = O(L^{D-1} \ln L)$  ( $L$ : length scale of  $A$ )
  - ▶ Recently, 1D solvable spin chain model which exhibit extensive entanglement entropy have been discussed.
    - ▶ Beyond logarithmic violation:  $S_A \propto \sqrt{(\text{volume of } A)}$   
[Movassagh, Shor 2014], [Salberger, Korepin 2016]
- Counterexamples of the belief!**

Introduction

**Motzkin spin model**

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

Rényi entropy

Rényi entropy of Motzkin model

Summary and discussion

# Motzkin spin model 1

[Bravyi et al 2012]

- ▶ 1D spin chain at sites  $i \in \{1, 2, \dots, 2n\}$
- ▶ Spin-1 state at each site can be regarded as up, down and flat steps;

$$|u\rangle \Leftrightarrow \nearrow, \quad |d\rangle \Leftrightarrow \searrow, \quad |0\rangle \Leftrightarrow \longrightarrow$$


# Motzkin spin model 1

[Bravyi et al 2012]

- ▶ 1D spin chain at sites  $i \in \{1, 2, \dots, 2n\}$
- ▶ Spin-1 state at each site can be regarded as up, down and flat steps;

$$|u\rangle \Leftrightarrow \nearrow, \quad |d\rangle \Leftrightarrow \searrow, \quad |0\rangle \Leftrightarrow \longrightarrow$$

- ▶ Each spin configuration  $\Leftrightarrow$  length- $2n$  walk in  $(x, y)$  plane  
Example)



## Motzkin spin model 2

[Bravyi et al 2012]

Hamiltonian:  $H_{\text{Motzkin}} = H_{\text{bulk}} + H_{\text{bdy}}$

► Bulk part:  $H_{\text{bulk}} = \sum_{j=1}^{2n-1} \Pi_{j,j+1}$ ,

$$\Pi_{j,j+1} = |D\rangle_{j,j+1}\langle D| + |U\rangle_{j,j+1}\langle U| + |F\rangle_{j,j+1}\langle F|$$

(local interactions) with

$$|D\rangle \equiv \frac{1}{\sqrt{2}} (|0, d\rangle - |d, 0\rangle),$$

$$|U\rangle \equiv \frac{1}{\sqrt{2}} (|0, u\rangle - |u, 0\rangle),$$

$$|F\rangle \equiv \frac{1}{\sqrt{2}} (|0, 0\rangle - |u, d\rangle).$$

## Motzkin spin model 2

[Bravyi et al 2012]

Hamiltonian:  $H_{\text{Motzkin}} = H_{\text{bulk}} + H_{\text{bdy}}$

► Bulk part:  $H_{\text{bulk}} = \sum_{j=1}^{2n-1} \Pi_{j,j+1}$ ,


$$\Pi_{j,j+1} = |D\rangle_{j,j+1}\langle D| + |U\rangle_{j,j+1}\langle U| + |F\rangle_{j,j+1}\langle F|$$

(local interactions) with

$$|D\rangle \equiv \frac{1}{\sqrt{2}} (|0, d\rangle - |d, 0\rangle),$$

$$|U\rangle \equiv \frac{1}{\sqrt{2}} (|0, u\rangle - |u, 0\rangle),$$

$$|F\rangle \equiv \frac{1}{\sqrt{2}} (|0, 0\rangle - |u, d\rangle).$$



## Motzkin spin model 2

[Bravyi et al 2012]

Hamiltonian:  $H_{\text{Motzkin}} = H_{\text{bulk}} + H_{\text{bdy}}$

► Bulk part:  $H_{\text{bulk}} = \sum_{j=1}^{2n-1} \Pi_{j,j+1}$ ,


$$\Pi_{j,j+1} = |D\rangle_{j,j+1}\langle D| + |U\rangle_{j,j+1}\langle U| + |F\rangle_{j,j+1}\langle F|$$

(local interactions) with

$$|D\rangle \equiv \frac{1}{\sqrt{2}} (|0, d\rangle - |d, 0\rangle),$$

$$|U\rangle \equiv \frac{1}{\sqrt{2}} (|0, u\rangle - |u, 0\rangle),$$

$$|F\rangle \equiv \frac{1}{\sqrt{2}} (|0, 0\rangle - |u, d\rangle).$$



## Motzkin spin model 2

[Bravyi et al 2012]

Hamiltonian:  $H_{\text{Motzkin}} = H_{\text{bulk}} + H_{\text{bdy}}$

► Bulk part:  $H_{\text{bulk}} = \sum_{j=1}^{2n-1} \Pi_{j,j+1}$ ,


$$\Pi_{j,j+1} = |D\rangle_{j,j+1}\langle D| + |U\rangle_{j,j+1}\langle U| + |F\rangle_{j,j+1}\langle F|$$

(local interactions) with

$$|D\rangle \equiv \frac{1}{\sqrt{2}} (|0, d\rangle - |d, 0\rangle),$$

$$|U\rangle \equiv \frac{1}{\sqrt{2}} (|0, u\rangle - |u, 0\rangle),$$

$$|F\rangle \equiv \frac{1}{\sqrt{2}} (|0, 0\rangle - |u, d\rangle).$$



“gauge equivalence”.



# Motzkin spin model 3

[Bravyi et al 2012]

Hamiltonian:  $H_{\text{Motzkin}} = H_{\text{bulk}} + H_{\text{bdy}}$

▶ Boundary part:  $H_{\text{bdy}} = |d\rangle_1 \langle d| + |u\rangle_{2n} \langle u|$



Hamiltonian:  $H_{Motzkin} = H_{bulk} + H_{bdy}$

- ▶ Boundary part:  $H_{bdy} = |d\rangle_1 \langle d| + |u\rangle_{2n} \langle u|$



- ▶  $H_{Motzkin}$  is the sum of projection operators.  
⇒ Positive semi-definite spectrum
- ▶ We find the unique zero-energy ground state.

Hamiltonian:  $H_{Motzkin} = H_{bulk} + H_{bdy}$

- ▶ Boundary part:  $H_{bdy} = |d\rangle_1 \langle d| + |u\rangle_{2n} \langle u|$



- ▶  $H_{Motzkin}$  is the sum of projection operators.  
⇒ Positive semi-definite spectrum
- ▶ We find the unique zero-energy ground state.
  - ▶ Each projector in  $H_{Motzkin}$  annihilates the zero-energy state.  
⇒ Frustration free
- ▶ The ground state corresponds to random walks starting at  $(0,0)$  and ending at  $(2n,0)$  restricted to the region  $y \geq 0$  (Motzkin Walks (MWs)).

## Motzkin spin model 4

[Bravyi et al 2012]

In terms of  $S = 1$  spin matrices

$$S_z = \begin{pmatrix} 1 & & \\ & 0 & \\ & & -1 \end{pmatrix}, \quad S_{\pm} \equiv \frac{1}{\sqrt{2}}(S_x \pm iS_y) = \begin{pmatrix} & 1 & \\ & & \\ & & \end{pmatrix}, \begin{pmatrix} 1 & & \\ & & \\ & & 1 \end{pmatrix},$$


$$H_{bulk} = \frac{1}{2} \sum_{j=1}^{2n-1} \left[ 1_j 1_{j+1} - \frac{1}{4} S_{zj} S_{zj+1} - \frac{1}{4} S_{zj}^2 S_{zj+1} + \frac{1}{4} S_{zj} S_{zj+1}^2 \right. \\ \left. - \frac{3}{4} S_{zj}^2 S_{zj+1}^2 + S_{+j} (S_z S_-)_{j+1} + S_{-j} (S_+ S_z)_{j+1} - (S_- S_z)_j S_{+j+1} \right. \\ \left. - (S_z S_+)_j S_{-j+1} - (S_- S_z)_j (S_+ S_z)_{j+1} - (S_z S_+)_j (S_z S_-)_{j+1} \right], \\ H_{bdy} = \frac{1}{2} (S_z^2 - S_z)_1 + \frac{1}{2} (S_z^2 + S_z)_{2n}$$

Quartic spin interactions

# Motzkin spin model 5

[Bravyi et al 2012]

Example)  $2n = 4$  case,  
MWs:



Ground state:

$$|P_4\rangle = \frac{1}{\sqrt{9}} [ |0000\rangle + |ud00\rangle + |0ud0\rangle + |00ud\rangle \\ + |u0d0\rangle + |0u0d\rangle + |u00d\rangle + |udud\rangle \\ + |uudd\rangle ].$$

# Motzkin spin model 6

[Bravyi et al 2012]

## Note


Forbidden paths for the ground state

1. Path entering  $y < 0$  region



Forbidden by  $H_{bdy}$

2. Path ending at nonzero height



Forbidden by  $H_{bdy}$

## Motzkin spin model 7

[Bravyi et al 2012]

Entanglement entropy of the subsystem  $A = \{1, 2, \dots, n\}$ :

- ▶ Normalization factor of the ground state  $|P_{2n}\rangle$  is given by the number of MWs of length  $2n$ :  $M_{2n} = \sum_{k=0}^n C_k \binom{2n}{2k}$ .

$$C_k = \frac{1}{k+1} \binom{2k}{k}: \text{Catalan number}$$

# Motzkin spin model 7

[Bravyi et al 2012]

Entanglement entropy of the subsystem  $A = \{1, 2, \dots, n\}$ :

- ▶ Normalization factor of the ground state  $|P_{2n}\rangle$  is given by the number of MWs of length  $2n$ :  $M_{2n} = \sum_{k=0}^n C_k \binom{2n}{2k}$ .

$$C_k = \frac{1}{k+1} \binom{2k}{k}: \text{Catalan number}$$

- ▶ Consider to trace out the density matrix  $\rho = |P_{2n}\rangle\langle P_{2n}|$  w.r.t. the subsystem  $B = \{n+1, \dots, 2n\}$ .

Schmidt decomposition:

$$|P_{2n}\rangle = \sum_{h \geq 0} \sqrt{p_{n,n}^{(h)}} |P_n^{(0 \rightarrow h)}\rangle \otimes |P_n^{(h \rightarrow 0)}\rangle$$

$$\text{with } p_{n,n}^{(h)} \equiv \frac{\binom{M_n^{(h)}}{M_{2n}}^2}{M_{2n}}.$$

↑  
Paths from  $(0, 0)$  to  $(n, h)$



# Motzkin spin model 8

[Bravyi et al 2012]

- ▶  $M_n^{(h)}$  is the number of paths in  $P_n^{(0 \rightarrow h)}$ .

For  $n \rightarrow \infty$ ,

Gaussian distribution

$$p_{n,n}^{(h)} \sim \frac{3\sqrt{6}}{\sqrt{\pi}} \frac{(h+1)^2}{n^{3/2}} e^{-\frac{3}{2} \frac{(h+1)^2}{n}} \times [1 + O(1/n)].$$

- ▶ Reduced density matrix

$$\rho_A = \text{Tr}_B \rho = \sum_{h \geq 0} p_{n,n}^{(h)} \left| P_n^{(0 \rightarrow h)} \right\rangle \left\langle P_n^{(0 \rightarrow h)} \right|$$

- ▶ Entanglement entropy

$$\begin{aligned} S_A &= - \sum_{h \geq 0} p_{n,n}^{(h)} \ln p_{n,n}^{(h)} \\ &= \frac{1}{2} \ln n + \frac{1}{2} \ln \frac{2\pi}{3} + \gamma - \frac{1}{2} \end{aligned} \quad (\gamma: \text{Euler constant})$$

up to terms vanishing as  $n \rightarrow \infty$ .

## Notes

- ▶ The system is critical (gapless).  
 $S_A$  is similar to the  $(1 + 1)$ -dimensional CFT with  $c = 3/2$ .

## Notes

- ▶ The system is critical (gapless).  
 $S_A$  is similar to the  $(1 + 1)$ -dimensional CFT with  $c = 3/2$ .
- ▶ But, gap scales as  $O(1/n^z)$  with  $z \geq 2$ .  
The system cannot be described by relativistic CFT.  
Lifshitz type ?  
Different  $z$  depending on excited states (Multiple dynamics)?

[Chen, Fradkin, Witczak-Krempa 2017]

## Notes

- ▶ The system is critical (gapless).  
 $S_A$  is similar to the  $(1 + 1)$ -dimensional CFT with  $c = 3/2$ .
- ▶ But, gap scales as  $O(1/n^z)$  with  $z \geq 2$ .  
The system cannot be described by relativistic CFT.  
Lifshitz type ?  
Different  $z$  depending on excited states (Multiple dynamics)?  
[Chen, Fradkin, Witczak-Krempa 2017]
- ▶ Excitations have not been much investigated.

Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

Rényi entropy

Rényi entropy of Motzkin model

Summary and discussion

# Colored Motzkin spin model 1

[Movassagh, Shor 2014]

- ▶ Introducing color d.o.f.  $k = 1, 2, \dots, s$  to up and down spins as

$$|u^k\rangle \Leftrightarrow \begin{array}{c} \nearrow \\ k \end{array}, \quad |d^k\rangle \Leftrightarrow \begin{array}{c} \searrow \\ k \end{array}, \quad |0\rangle \Leftrightarrow \longrightarrow$$

Color d.o.f. decorated to Motzkin Walks

# Colored Motzkin spin model 1

[Movassagh, Shor 2014]

- ▶ Introducing color d.o.f.  $k = 1, 2, \dots, s$  to up and down spins as

$$|u^k\rangle \Leftrightarrow \begin{array}{c} \nearrow k \\ \text{red arrow} \end{array}, \quad |d^k\rangle \Leftrightarrow \begin{array}{c} \searrow k \\ \text{red arrow} \end{array}, \quad |0\rangle \Leftrightarrow \longrightarrow$$

Color d.o.f. decorated to Motzkin Walks

- ▶ Hamiltonian  $H_{cMotzkin} = H_{bulk} + H_{bdy}$

- ▶ Bulk part consisting of **local interactions**:

$$H_{bulk} = \sum_{j=1}^{2n-1} (\Pi_{j,j+1} + \Pi_{j,j+1}^{cross}),$$

$$\Pi_{j,j+1} = \sum_{k=1}^s \left[ |D^k\rangle_{j,j+1} \langle D^k| + |U^k\rangle_{j,j+1} \langle U^k| + |F^k\rangle_{j,j+1} \langle F^k| \right]$$

with

$$|D^k\rangle \equiv \frac{1}{\sqrt{2}} \left( |0, d^k\rangle - |d^k, 0\rangle \right),$$

$$|U^k\rangle \equiv \frac{1}{\sqrt{2}} \left( |0, u^k\rangle - |u^k, 0\rangle \right),$$

$$|F^k\rangle \equiv \frac{1}{\sqrt{2}} \left( |0, 0\rangle - |u^k, d^k\rangle \right),$$

and

$$\Pi_{j,j+1}^{\text{cross}} = \sum_{k \neq k'} |u^k, d^{k'}\rangle_{j,j+1} \langle u^k, d^{k'}|.$$

⇒ Colors should be matched in up and down pairs.

► Boundary part

$$H_{\text{bdy}} = \sum_{k=1}^s \left( |d^k\rangle_1 \langle d^k| + |u^k\rangle_{2n} \langle u^k| \right).$$



## Colored Motzkin spin model 3


[Movassagh, Shor 2014]

- ▶ Still unique ground state with zero energy

# Colored Motzkin spin model 3

[Movassagh, Shor 2014]

- ▶ Still unique ground state with zero energy
- ▶ Example)  $2n = 4$  case,



$$\begin{aligned}
 |P_4\rangle = & \frac{1}{\sqrt{1 + 6s + 2s^2}} \left[ |0000\rangle + \sum_{k=1}^s \left\{ |u^k d^k 00\rangle + \dots + |u^k 00 d^k\rangle \right\} \right. \\
 & \left. + \sum_{k,k'=1}^s \left\{ |u^k d^k u^{k'} d^{k'}\rangle + |u^k u^{k'} d^{k'} d^k\rangle \right\} \right].
 \end{aligned}$$

## Entanglement entropy

- ▶ Paths from  $(0, 0)$  to  $(n, h)$ ,  $P_n^{(0 \rightarrow h)}$ , have  $h$  unmatched up steps.

Let  $\tilde{P}_n^{(0 \rightarrow h)}(\{\kappa_m\})$  be paths with the colors of unmatched up steps frozen.

(unmatched up from height  $(m - 1)$  to  $m$ )  $\rightarrow u^{\kappa_m}$

- ▶ Similarly,

$$P_n^{(h \rightarrow 0)} \rightarrow \tilde{P}_n^{(h \rightarrow 0)}(\{\kappa_m\}),$$

(unmatched down from height  $m$  to  $(m - 1)$ )  $\rightarrow d^{\kappa_m}$ .


- ▶ The numbers satisfy  $M_n^{(h)} = s^h \tilde{M}_n^{(h)}$ .

# Colored Motzkin spin model 5

[Movassagh, Shor 2014]

## Example

$2n = 8$  case,  $h = 2$



- ▶ Schmidt decomposition

$$\begin{aligned}
 |P_{2n}\rangle &= \sum_{h \geq 0} \sum_{\kappa_1=1}^s \cdots \sum_{\kappa_h=1}^s \sqrt{\rho_{n,n}^{(h)}} \\
 &\quad \times \left| \tilde{P}_n^{(0 \rightarrow h)}(\{\kappa_m\}) \right\rangle \otimes \left| \tilde{P}_n^{(h \rightarrow 0)}(\{\kappa_m\}) \right\rangle
 \end{aligned}$$

with

$$\rho_{n,n}^{(h)} = \frac{\left( \tilde{M}_n^{(h)} \right)^2}{M_{2n}}.$$

- ▶ Reduced density matrix

$$\begin{aligned}
 \rho_A &= \sum_{h \geq 0} \sum_{\kappa_1=1}^s \cdots \sum_{\kappa_h=1}^s \rho_{n,n}^{(h)} \\
 &\quad \times \left| \tilde{P}_n^{(0 \rightarrow h)}(\{\kappa_m\}) \right\rangle \left\langle \tilde{P}_n^{(0 \rightarrow h)}(\{\kappa_m\}) \right|.
 \end{aligned}$$

- ▶ For  $n \rightarrow \infty$ ,

$$p_{n,n}^{(h)} \sim \frac{\sqrt{2} s^{-h}}{\sqrt{\pi} (\sigma n)^{3/2}} (h+1)^2 e^{-\frac{(h+1)^2}{2\sigma n}} \times [1 + O(1/n)]$$

with  $\sigma \equiv \frac{\sqrt{s}}{2\sqrt{s+1}}$ .

Note: Effectively  $h \lesssim O(\sqrt{n})$ .

- ▶ Entanglement entropy

$$S_A = - \sum_{h \geq 0} s^h p_{n,n}^{(h)} \ln p_{n,n}^{(h)}$$

- ▶ For  $n \rightarrow \infty$ ,

$$p_{n,n}^{(h)} \sim \frac{\sqrt{2} s^{-h}}{\sqrt{\pi} (\sigma n)^{3/2}} (h+1)^2 e^{-\frac{(h+1)^2}{2\sigma n}} \times [1 + O(1/n)]$$

with  $\sigma \equiv \frac{\sqrt{s}}{2\sqrt{s+1}}$ .

Note: Effectively  $h \lesssim O(\sqrt{n})$ .

- ▶ Entanglement entropy

$$\begin{aligned} S_A &= - \sum_{h \geq 0} s^h p_{n,n}^{(h)} \ln p_{n,n}^{(h)} \\ &= (2 \ln s) \sqrt{\frac{2\sigma n}{\pi}} + \frac{1}{2} \ln n + \frac{1}{2} \ln(2\pi\sigma) + \gamma - \frac{1}{2} - \ln s \end{aligned}$$

up to terms vanishing as  $n \rightarrow \infty$ .

Grows as  $\sqrt{n}$ .

## Comments

- ▶ Matching color  $\Rightarrow s^{-h}$  factor in  $p_{n,n}^{(h)}$   
 $\Rightarrow$  crucial to  $O(\sqrt{n})$  behavior in  $S_A$




# Colored Motzkin spin model 8

[Movassagh, Shor 2014]

## Comments

- ▶ Matching color  $\Rightarrow s^{-h}$  factor in  $p_{n,n}^{(h)}$   
 $\Rightarrow$  crucial to  $O(\sqrt{n})$  behavior in  $S_A$
- ▶ Typical configurations:




+ (equivalence moves).

# Colored Motzkin spin model 8

[Movassagh, Shor 2014]

## Comments

- ▶ Matching color  $\Rightarrow s^{-h}$  factor in  $p_{n,n}^{(h)}$   
 $\Rightarrow$  crucial to  $O(\sqrt{n})$  behavior in  $S_A$
- ▶ Typical configurations:



+ (equivalence moves).

- ▶ For spin 1/2 chain (**only up and down**), the model in which similar behavior exhibits in colored as well as uncolored cases has been constructed. (**Fredkin model**) [Salberger, Korepin 2016]

# Colored Motzkin spin model 9

[Movassagh, Shor 2014]

► Correlation functions

[Dell'Anna et al, 2016]

$$\langle S_{z,1} S_{z,2n} \rangle_{\text{connected}} \rightarrow -0.034... \times \frac{s^3 - s}{6} \neq 0 \quad (n \rightarrow \infty)$$

⇒ Violation of cluster decomposition property for  $s > 1$   
(Strong correlation due to color matching)

# Colored Motzkin spin model 9

[Movassagh, Shor 2014]

- ▶ Correlation functions

[Dell'Anna et al, 2016]

$$\langle S_{z,1} S_{z,2n} \rangle_{\text{connected}} \rightarrow -0.034\dots \times \frac{s^3 - s}{6} \neq 0 \quad (n \rightarrow \infty)$$

⇒ Violation of cluster decomposition property for  $s > 1$   
(Strong correlation due to color matching)

- ▶ Deformation of models to achieve the volume law behavior  
( $S_A \propto n$ )

Weighted Motzkin/Dyck walks

[Zhang et al, Salberger et al 2016]

Introduction

Motzkin spin model

Colored Motzkin model

**SIS Motzkin model**

Colored SIS Motzkin model

Rényi entropy

Rényi entropy of Motzkin model

Summary and discussion

# Symmetric Inverse Semigroups (SISs)

- ▶ Inverse Semigroup ( $\subset$  Semigroup):  
An unique inverse exists for every element.  
But, no unique identity (partial identities).

# Symmetric Inverse Semigroups (SISs)

- ▶ Inverse Semigroup ( $\subset$  Semigroup):  
An unique inverse exists for every element.  
But, no unique identity (partial identities).

- ▶ SIS ( $\subset$  Semigroup):

Semigroup version of the symmetric group  $S_k$

$$S_p^k \quad (p = 1, \dots, k)$$

# Symmetric Inverse Semigroups (SISs)

- ▶ Inverse Semigroup ( $\subset$  Semigroup):  
An unique inverse exists for every element.  
But, no unique identity (partial identities).

- ▶ SIS ( $\subset$  Semigroup):

Semigroup version of the symmetric group  $S_k$

$$S_p^k \quad (p = 1, \dots, k)$$

- ▶  $x_{a,b} \in S_1^k$  maps  $a$  to  $b$ . ( $a, b \in \{1, \dots, k\}$ )

Product rule:  $x_{a,b} * x_{c,d} = \delta_{b,c} x_{a,d}$

$$x_{1,2} * x_{2,1} = x_{1,1}, \quad x_{2,1} * x_{1,2} = x_{2,2}$$



(partial identities)

$$(x_{1,2})^{-1} = x_{2,1}$$

(unique inverse)



# Symmetric Inverse Semigroups (SISs)

- ▶ **Inverse Semigroup** ( $\subset$  Semigroup):  
An unique inverse exists for every element.  
But, no unique identity (partial identities).

- ▶ **SIS** ( $\subset$  Semigroup):

Semigroup version of the symmetric group  $S_k$

$$S_p^k \quad (p = 1, \dots, k)$$

- ▶  $x_{a,b} \in S_1^k$  maps  $a$  to  $b$ . ( $a, b \in \{1, \dots, k\}$ )

Product rule:  $x_{a,b} * x_{c,d} = \delta_{b,c} x_{a,d}$

$$x_{1,2} * x_{2,1} = x_{1,1}, \quad x_{2,1} * x_{1,2} = x_{2,2}$$

(partial identities)

$$(x_{1,2})^{-1} = x_{2,1} \quad (\text{unique inverse})$$

- ▶  $x_{a_1, a_2}; b_1, b_2 \in S_2^k$  etc, ...

# Symmetric Inverse Semigroups (SISs)

- ▶ **Inverse Semigroup** ( $\subset$  Semigroup):  
An unique inverse exists for every element.  
But, no unique identity (partial identities).

- ▶ **SIS** ( $\subset$  Semigroup):

Semigroup version of the symmetric group  $S_k$

$$S_p^k \quad (p = 1, \dots, k)$$

- ▶  $x_{a,b} \in S_1^k$  maps  $a$  to  $b$ . ( $a, b \in \{1, \dots, k\}$ )

Product rule:  $x_{a,b} * x_{c,d} = \delta_{b,c} x_{a,d}$

$$x_{1,2} * x_{2,1} = x_{1,1}, \quad x_{2,1} * x_{1,2} = x_{2,2}$$

(partial identities)

$$(x_{1,2})^{-1} = x_{2,1} \quad (\text{unique inverse})$$




- ▶  $x_{a_1, a_2}; b_1, b_2 \in S_2^k$  etc, ...

$$S_k^k \equiv S_k$$


# SIS Motzkin model 1


[Sugino, Padmanabhan 2017]


- ▶ Change the spin d.o.f. as  $|x_{a,b}\rangle$  with  $a, b \in \{1, 2, \dots, k\}$ .

- ▶  $a < b$  case: 'up'  $\Leftrightarrow$  
- ▶  $a > b$  case: 'down'  $\Leftrightarrow$  
- ▶  $a = b$  case: 'flat'  $\Leftrightarrow$  

- ▶ Change the spin d.o.f. as  $|x_{a,b}\rangle$  with  $a, b \in \{1, 2, \dots, k\}$ .

- ▶  $a < b$  case: 'up'  $\Leftrightarrow$  

- $a > b$  case: 'down'  $\Leftrightarrow$  

- $a = b$  case: 'flat'  $\Leftrightarrow$  

- ▶ We regard the configuration of adjacent sites  $|x_{a,b}\rangle_j |x_{c,d}\rangle_{j+1}$  as a connected path for  $b = c$ .


c.f.) Analogous to the product rule of Symmetric Inverse Semigroup ( $\mathcal{S}_1^k$ ):

$$x_{a,b} * x_{c,d} = \delta_{b,c} x_{a,d}$$

$a, b$ : semigroup indices

- ▶ Inner product:  $\langle x_{a,b} | x_{c,d} \rangle = \delta_{a,c} \delta_{b,d}$
- ▶ Let us consider the  $k = 3$  case.

- ▶ Maximum height is lower than the original Motzkin case.




# SIS Motzkin model 3

[Sugino, Padmanabhan 2017]

Hamiltonian  $H_{S31Motzkin} = H_{bulk} + H_{bulk,disc} + H_{bdy}$

►  $H_{bulk}$ : **local interactions** corresponding to the following moves:



- ▶  $H_{bulk,disc}$  lifts disconnected paths to excited states.

$\Pi^{|\psi\rangle}$ : projector to  $|\psi\rangle$

$$H_{bulk,disc} = \sum_{j=1}^{2n-1} \sum_{a,b,c,d=1; b \neq c}^3 \Pi^{|(x_{a,b})_j, (x_{c,d})_{j+1}\rangle}$$

- ▶  $H_{bulk,disc}$  lifts disconnected paths to excited states.

$\Pi^{|\psi\rangle}$ : projector to  $|\psi\rangle$

$$H_{bulk,disc} = \sum_{j=1}^{2n-1} \sum_{a,b,c,d=1; b \neq c}^3 \Pi^{|(x_{a,b})_j, (x_{c,d})_{j+1}\rangle}$$

- ▶

$$H_{bdy} = \sum_{a>b} \Pi^{|(x_{a,b})_1\rangle} + \sum_{a<b} \Pi^{|(x_{a,b})_{2n}\rangle} \\ + \Pi^{|(x_{1,3})_1, (x_{3,2})_2, (x_{2,1})_3\rangle} + \Pi^{|(x_{1,2})_{2n-2}, (x_{2,3})_{2n-1}, (x_{3,1})_{2n}\rangle}$$

The last 2 terms have no analog to the original Motzkin model.



## SIS Motzkin model 5

[Sugino, Padmanabhan 2017]

- ▶ Ground states correspond to connected paths starting at  $(0,0)$ , ending at  $(2n,0)$  and not entering  $y < 0$ .  $\mathcal{S}_1^3$  MWs

## SIS Motzkin model 5

[Sugino, Padmanabhan 2017]

- ▶ Ground states correspond to connected paths starting at  $(0,0)$ , ending at  $(2n,0)$  and not entering  $y < 0$ .  $\mathcal{S}_1^3$  MWs
- ▶ The ground states have 5 fold degeneracy according to the initial and final semigroup indices:  
 $(1,1)$ ,  $(1,2)$ ,  $(2,1)$ ,  $(2,2)$  and  $(3,3)$  sectors  
The  $(3,3)$  sector is trivial, consisting of only one path:

$$x_{3,3} x_{3,3} \cdots x_{3,3}$$

## SIS Motzkin model 5

[Sugino, Padmanabhan 2017]

- ▶ Ground states correspond to connected paths starting at  $(0,0)$ , ending at  $(2n,0)$  and not entering  $y < 0$ .  $S_1^3$  MWs
- ▶ The ground states have 5 fold degeneracy according to the initial and final semigroup indices:  
 $(1,1)$ ,  $(1,2)$ ,  $(2,1)$ ,  $(2,2)$  and  $(3,3)$  sectors

The  $(3,3)$  sector is trivial, consisting of only one path:

$$x_{3,3}x_{3,3} \cdots x_{3,3}$$

- ▶ The number of paths can be obtained by recursion relations. For length- $n$  paths from the semigroup index  $a$  to  $b$  ( $P_{n,a \rightarrow b}$ ),

$$\begin{aligned} P_{n,1 \rightarrow 1} &= x_{1,1}P_{n-1,1 \rightarrow 1} + x_{1,2} \sum_{i=1}^{n-2} P_{i,2 \rightarrow 2} x_{2,1}P_{n-2-i,1 \rightarrow 1} \\ &\quad + x_{1,3} \sum_{i=1}^{n-2} P_{i,3 \rightarrow 3} x_{3,1}P_{n-2-i,1 \rightarrow 1} \\ &\quad + x_{1,3} \sum_{i=1}^{n-2} P_{i,3 \rightarrow 3} x_{3,2}P_{n-2-i,2 \rightarrow 1}, \quad \text{etc.} \end{aligned}$$

## Result

- ▶ The entanglement entropies  $S_{A,1\rightarrow 1}$ ,  $S_{A,1\rightarrow 2}$ ,  $S_{A,2\rightarrow 1}$  and  $S_{A,2\rightarrow 2}$  take the same form as in the case of the Motzkin model.

Logarithmic violation of the area law


- ▶ The form of  $p_n^{(h)} \sim \frac{(h+1)^2}{n^{3/2}} e^{-(\text{const.})\frac{(h+1)^2}{n}}$  is universal.
- ▶  $S_{A,3\rightarrow 3} = 0$ .

# SIS Motzkin model 7

## Localization

[Padmanabhan, F.S., Korepin 2018]

- ▶ There are excited states corresponding to disconnected paths.  
Example) One such path in  $2n = 6$  case,




# SIS Motzkin model 7

## Localization

[Padmanabhan, F.S., Korepin 2018]

- ▶ There are excited states corresponding to disconnected paths.  
Example) One such path in  $2n = 6$  case,



Corresponding excited state:  $|P_{3,1 \rightarrow 1}\rangle \otimes |P_{3,2 \rightarrow 1}^{(1 \rightarrow 0)}\rangle$

Each connected component has no entanglement with other components.


“2nd quantization” of paths

# SIS Motzkin model 7

## Localization

[Padmanabhan, F.S., Korepin 2018]

- ▶ There are excited states corresponding to disconnected paths.  
Example) One such path in  $2n = 6$  case,



Corresponding excited state:  $|P_{3,1 \rightarrow 1}\rangle \otimes |P_{3,2 \rightarrow 1}^{(1 \rightarrow 0)}\rangle$

Each connected component has no entanglement with other components.

“2nd quantization” of paths

$\Rightarrow$  2pt connected correlation functions of local operators belonging to separate connected components vanish.

$\Rightarrow$  Localization!

Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

**Colored SIS Motzkin model**

Rényi entropy

Rényi entropy of Motzkin model

Summary and discussion



The SIS  $\mathcal{S}_2^3$

- ▶ 18 elements  $x_{ab,cd}$  with  $ab \in \{12, 23, 31\}$  and  $cd \in \{12, 23, 31, 21, 32, 13\}$  satisfying

$$x_{ab,cd} * x_{ef,gh} = \delta_{c,e} \delta_{d,f} x_{ab,gh} + \delta_{c,f} \delta_{d,e} x_{ab,hg}.$$

- ▶ can be regarded as 2 sets of  $\mathcal{S}_1^3$ . ⇒ color d.o.f.

The SIS  $\mathcal{S}_2^3$

- ▶ 18 elements  $x_{ab,cd}$  with  $ab \in \{12, 23, 31\}$  and  $cd \in \{12, 23, 31, 21, 32, 13\}$  satisfying

$$x_{ab,cd} * x_{ef,gh} = \delta_{c,e}\delta_{d,f} x_{ab,gh} + \delta_{c,f}\delta_{d,e} x_{ab,hg}.$$

- ▶ can be regarded as 2 sets of  $\mathcal{S}_1^3$ . ⇒ color d.o.f.
- ▶ Spin variables:  $x_{a,b}^s$  ( $s = 1, 2$ ) ( $a, b = 1, 2, 3$ )
- ▶ The new moves (C moves) introduced to the Hamiltonian.

$$\overset{1}{\longrightarrow} a \sim a \overset{2}{\longrightarrow}$$

# Colored SIS Motzkin model 2

[Sugino, Padmanabhan 2017]

Hamiltonian:  $H_{cS31Motzkin} = H_{bulk} + H_{bulk,disc} + H_{bdy}$


- ▶ In  $H_{bulk}$ , (Down), (Up) and (Flat) are essentially the same as before.



# Colored SIS Motzkin model 3

[Sugino, Padmanabhan 2017]

- ▶ Wedge move:



- ▶

$$(Cross)_{j,j+1} = \sum_{b>a,c} \left[ \prod |(x_{a,b}^1)_j, (x_{b,c}^2)_{j+1}\rangle + \prod |(x_{a,b}^2)_j, (x_{b,c}^1)_{j+1}\rangle \right]$$

forbids unmatched up and down steps in ground states.

$\Downarrow$

$$H_{bulk} = \mu \sum_{j=1}^{2n} C_j + \sum_{j=1}^{2n-1} [(Down)_{j,j+1} + (Up)_{j,j+1} \\ + (Flat)_{j,j+1} + (Wedge)_{j,j+1} + (Cross)_{j,j+1}]$$



$$H_{bulk,disc} = \sum_{j=1}^{2n-1} \sum_{a,b,c,d=1; b \neq c}^3 \sum_{s,t=1}^2 \prod |(x_{a,b}^s)_j, (x_{c,d}^t)_{j+1}\rangle$$



$$\begin{aligned}
 H_{bdy} = & \sum_{a>b} \sum_{s=1}^2 \prod |(x_{a,b}^s)_1\rangle + \sum_{a<b} \sum_{s=1}^2 \prod |(x_{a,b}^s)_{2n}\rangle \\
 & + \sum_{s,t=1}^2 \prod |(x_{1,3}^s)_1, (x_{3,2}^s)_2, (x_{2,1}^t)_3\rangle \\
 & + \sum_{s,t=1}^2 \prod |(x_{1,2}^s)_{2n-2}, (x_{2,3}^t)_{2n-1}, (x_{3,1}^t)_{2n}\rangle
 \end{aligned}$$

- ▶ 5 ground states of (1, 1), (1, 2), (2, 1), (2, 2), (3, 3) sectors
- ▶ Quantum phase transition between  $\mu > 0$  and  $\mu = 0$  in the 4 sectors except (3, 3).
  - ▶ For  $\mu > 0$ ,

$$S_A = (2 \ln 2) \sqrt{\frac{2\sigma n}{\pi}} + \frac{1}{2} \ln n + \frac{1}{2} \ln(2\pi\sigma) + \gamma - \frac{1}{2} + \ln \frac{3}{2^{1/3}}$$

with  $\sigma \equiv \frac{\sqrt{2}-1}{9\sqrt{2}}$ .

- ▶ For  $\mu = 0$ , colors 1 and 2 decouple.

$$S_A \propto \ln n.$$

Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

**Rényi entropy**

Rényi entropy of Motzkin model

Summary and discussion

# Rényi entropy

[Rényi, 1970]

- ▶ Rényi entropy has further importance than the von Neumann entanglement entropy:

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \text{Tr}_A \rho_A^\alpha \quad \text{with } \alpha > 0 \text{ and } \alpha \neq 1.$$



# Rényi entropy

[Rényi, 1970]

- ▶ Rényi entropy has further importance than the von Neumann entanglement entropy:

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \text{Tr}_A \rho_A^\alpha \quad \text{with } \alpha > 0 \text{ and } \alpha \neq 1.$$

- ▶ Generalization of the von Neumann entanglement entropy:  
 $\lim_{\alpha \rightarrow 1} S_{A,\alpha} = S_A$

# Rényi entropy

[Rényi, 1970]

- ▶ Rényi entropy has further importance than the von Neumann entanglement entropy:

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \text{Tr}_A \rho_A^\alpha \quad \text{with } \alpha > 0 \text{ and } \alpha \neq 1.$$

- ▶ Generalization of the von Neumann entanglement entropy:  
 $\lim_{\alpha \rightarrow 1} S_{A,\alpha} = S_A$
- ▶ Reconstructs the whole spectrum of the entanglement Hamiltonian  $H_{\text{ent},A} \equiv -\ln \rho_A$ .

# Rényi entropy

[Rényi, 1970]

- ▶ Rényi entropy has further importance than the von Neumann entanglement entropy:

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \text{Tr}_A \rho_A^\alpha \quad \text{with } \alpha > 0 \text{ and } \alpha \neq 1.$$

- ▶ Generalization of the von Neumann entanglement entropy:

$$\lim_{\alpha \rightarrow 1} S_{A,\alpha} = S_A$$

- ▶ Reconstructs the whole spectrum of the entanglement

Hamiltonian  $H_{\text{ent},A} \equiv -\ln \rho_A$ .

- ▶ For  $S_{A,\alpha}$  ( $0 < \alpha < 1$ ), the gapped systems in 1D is proven to obey the area law. [Huang, 2015]

- ▶ Rényi entropy has further importance than the von Neumann entanglement entropy:

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \text{Tr}_A \rho_A^\alpha \quad \text{with } \alpha > 0 \text{ and } \alpha \neq 1.$$

- ▶ Generalization of the von Neumann entanglement entropy:  
 $\lim_{\alpha \rightarrow 1} S_{A,\alpha} = S_A$
- ▶ Reconstructs the whole spectrum of the entanglement Hamiltonian  $H_{\text{ent},A} \equiv -\ln \rho_A$ .
- ▶ For  $S_{A,\alpha}$  ( $0 < \alpha < 1$ ), the gapped systems in 1D is proven to obey the area law. [Huang, 2015]

Here, I give a review of Motzkin spin chain and analytically compute its Rényi entropy of half-chain.

New phase transition found at  $\alpha = 1$ !

Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

Rényi entropy

**Rényi entropy of Motzkin model**

Summary and discussion

- ▶ What we compute is the asymptotic behavior of

$$S_{A, \alpha} = \frac{1}{1 - \alpha} \ln \sum_{h=0}^n s^h \left( p_{n,n}^{(h)} \right)^\alpha .$$

- ▶ What we compute is the asymptotic behavior of

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \sum_{h=0}^n s^h \left( p_{n,n}^{(h)} \right)^\alpha.$$

- ▶ For colorless case ( $s = 1$ ), we obtain

$$\begin{aligned} S_{A,\alpha} &= \frac{1}{2} \ln n + \frac{1}{1-\alpha} \ln \Gamma \left( \alpha + \frac{1}{2} \right) \\ &\quad - \frac{1}{2(1-\alpha)} \left\{ (1+2\alpha) \ln \alpha + \alpha \ln \frac{\pi}{24} + \ln 6 \right\} \end{aligned}$$

up to terms vanishing as  $n \rightarrow \infty$ .

- ▶ What we compute is the asymptotic behavior of

$$S_{A,\alpha} = \frac{1}{1-\alpha} \ln \sum_{h=0}^n s^h \left( p_{n,n}^{(h)} \right)^\alpha.$$

- ▶ For colorless case ( $s = 1$ ), we obtain

$$\begin{aligned} S_{A,\alpha} = & \frac{1}{2} \ln n + \frac{1}{1-\alpha} \ln \Gamma \left( \alpha + \frac{1}{2} \right) \\ & - \frac{1}{2(1-\alpha)} \left\{ (1+2\alpha) \ln \alpha + \alpha \ln \frac{\pi}{24} + \ln 6 \right\} \end{aligned}$$

up to terms vanishing as  $n \rightarrow \infty$ .

- ▶ Logarithmic growth
- ▶ Reduces to  $S_A$  in the  $\alpha \rightarrow 1$  limit.
- ▶ Consistent with half-chain case in the result in [Movassagh, 2017]



## Rényni entropy of Motzkin model 2

[F.S., Korepin, 2018]

Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .

## Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .

For  $0 < \alpha < 1$ , exponentially growing (colored case ( $s > 1$ )).

$\Rightarrow$  Saddle point value of the sum:  $h_* = O(n)$

## Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .  
For  $0 < \alpha < 1$ , exponentially growing (colored case ( $s > 1$ )).  
 $\Rightarrow$  Saddle point value of the sum:  $h_* = O(n)$
- ▶ Saddle point analysis for the sum leads to

$$S_{A,\alpha} = n \frac{2\alpha}{1-\alpha} \ln \left[ \sigma \left( s^{\frac{1-\alpha}{2\alpha}} + s^{-\frac{1-\alpha}{2\alpha}} + s^{-1/2} \right) \right] \\ + \frac{1+\alpha}{2(1-\alpha)} \ln n + C(s, \alpha)$$

with  $C(s, \alpha)$  being  $n$ -independent terms.

## Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .  
For  $0 < \alpha < 1$ , exponentially growing (colored case ( $s > 1$ )).  
 $\Rightarrow$  Saddle point value of the sum:  $h_* = O(n)$
- ▶ Saddle point analysis for the sum leads to

$$S_{A,\alpha} = n \frac{2\alpha}{1-\alpha} \ln \left[ \sigma \left( s^{\frac{1-\alpha}{2\alpha}} + s^{-\frac{1-\alpha}{2\alpha}} + s^{-1/2} \right) \right] \\ + \frac{1+\alpha}{2(1-\alpha)} \ln n + C(s, \alpha)$$

with  $C(s, \alpha)$  being  $n$ -independent terms.

- ▶ The saddle point value is  $h_* = n \frac{s^{\frac{1}{2\alpha}} - s^{1-\frac{1}{2\alpha}}}{s^{\frac{1}{2\alpha}} + s^{1-\frac{1}{2\alpha}} + 1} + O(n^0)$ .

## Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .

For  $0 < \alpha < 1$ , exponentially growing (colored case ( $s > 1$ )).

⇒ Saddle point value of the sum:  $h_* = O(n)$

- ▶ Saddle point analysis for the sum leads to

$$S_{A,\alpha} = n \frac{2\alpha}{1-\alpha} \ln \left[ \sigma \left( s^{\frac{1-\alpha}{2\alpha}} + s^{-\frac{1-\alpha}{2\alpha}} + s^{-1/2} \right) \right] \\ + \frac{1+\alpha}{2(1-\alpha)} \ln n + C(s, \alpha)$$

with  $C(s, \alpha)$  being  $n$ -independent terms.

- ▶ The saddle point value is  $h_* = n \frac{s^{\frac{1}{2\alpha}} - s^{1-\frac{1}{2\alpha}}}{s^{\frac{1}{2\alpha}} + s^{1-\frac{1}{2\alpha}} + 1} + O(n^0)$ .
- ▶ Linear growth in  $n$ .

## Colored case ( $s > 1$ )

- ▶ The summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  has a factor  $s^{(1-\alpha)h}$ .

For  $0 < \alpha < 1$ , exponentially growing (colored case ( $s > 1$ )).

⇒ Saddle point value of the sum:  $h_* = O(n)$

- ▶ Saddle point analysis for the sum leads to

$$S_{A,\alpha} = n \frac{2\alpha}{1-\alpha} \ln \left[ \sigma \left( s^{\frac{1-\alpha}{2\alpha}} + s^{-\frac{1-\alpha}{2\alpha}} + s^{-1/2} \right) \right] \\ + \frac{1+\alpha}{2(1-\alpha)} \ln n + C(s, \alpha)$$

with  $C(s, \alpha)$  being  $n$ -independent terms.

- ▶ The saddle point value is  $h_* = n \frac{s^{\frac{1}{2\alpha}} - s^{1-\frac{1}{2\alpha}}}{s^{\frac{1}{2\alpha}} + s^{1-\frac{1}{2\alpha}} + 1} + O(n^0)$ .
- ▶ Linear growth in  $n$ .
- ▶ Note:  $\alpha \rightarrow 1$  or  $s \rightarrow 1$  limit does not commute with the  $n \rightarrow \infty$  limit.

Rényni entropy for  $\alpha > 1$

- ▶ For  $\alpha > 1$ , the factor  $s^{(1-\alpha)h}$  in the summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  exponentially decays.

Rényni entropy for  $\alpha > 1$

- ▶ For  $\alpha > 1$ , the factor  $s^{(1-\alpha)h}$  in the summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  exponentially decays.  
 $\Rightarrow h \lesssim O\left(\frac{1}{(\alpha-1)\ln s}\right) = O(n^0)$  dominantly contributes to the sum.



## Rényi entropy for $\alpha > 1$

- ▶ For  $\alpha > 1$ , the factor  $s^{(1-\alpha)h}$  in the summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  exponentially decays.  
 $\Rightarrow h \lesssim O\left(\frac{1}{(\alpha-1)\ln s}\right) = O(n^0)$  dominantly contributes to the sum.
- ▶ The result:

$$S_{A,\alpha} = \frac{3\alpha}{2(\alpha-1)} \ln n + O(n^0).$$

## Rényi entropy for $\alpha > 1$

- ▶ For  $\alpha > 1$ , the factor  $s^{(1-\alpha)h}$  in the summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  exponentially decays.  
 $\Rightarrow h \lesssim O\left(\frac{1}{(\alpha-1)\ln s}\right) = O(n^0)$  dominantly contributes to the sum.
- ▶ The result:

$$S_{A,\alpha} = \frac{3\alpha}{2(\alpha-1)} \ln n + O(n^0).$$

- ▶ Logarithmic growth

## Rényi entropy for $\alpha > 1$

- ▶ For  $\alpha > 1$ , the factor  $s^{(1-\alpha)h}$  in the summand  $s^h \left( p_{n,n}^{(h)} \right)^\alpha$  exponentially decays.  
 $\Rightarrow h \lesssim O\left(\frac{1}{(\alpha-1)\ln s}\right) = O(n^0)$  dominantly contributes to the sum.
- ▶ The result:

$$S_{A,\alpha} = \frac{3\alpha}{2(\alpha-1)} \ln n + O(n^0).$$

- ▶ Logarithmic growth
- ▶  $\alpha \rightarrow 1$  or  $s \rightarrow 1$  limit does not commute with the  $n \rightarrow \infty$  limit.

# Rényni entropy of Motzkin model 4

[F.S., Korepin, 2018]

## Phase transition

- ▶  $S_{A^\alpha}$  grows as  $O(n)$  for  $0 < \alpha < 1$  while as  $O(\ln n)$  for  $\alpha > 1$ .

## Phase transition

- ▶  $S_{A_\alpha}$  grows as  $O(n)$  for  $0 < \alpha < 1$  while as  $O(\ln n)$  for  $\alpha > 1$ .  
⇒ Non-analytic behavior at  $\alpha = 1$  (Phase transition)

## Phase transition


- ▶  $S_{A^\alpha}$  grows as  $O(n)$  for  $0 < \alpha < 1$  while as  $O(\ln n)$  for  $\alpha > 1$ .  
⇒ Non-analytic behavior at  $\alpha = 1$  (Phase transition)
- ▶ In terms of the entanglement Hamiltonian,  
 $\text{Tr}_A \rho_A^\alpha = \text{Tr}_A e^{-\alpha H_{\text{ent}, A}}$        $\alpha$ : “inverse temperature”

## Phase transition

- ▶  $S_{A^\alpha}$  grows as  $O(n)$  for  $0 < \alpha < 1$  while as  $O(\ln n)$  for  $\alpha > 1$ .  
⇒ Non-analytic behavior at  $\alpha = 1$  (Phase transition)
- ▶ In terms of the entanglement Hamiltonian,  
$$\text{Tr}_A \rho_A^\alpha = \text{Tr}_A e^{-\alpha H_{\text{ent}, A}} \quad \alpha: \text{“inverse temperature”}$$
  - ▶  $0 < \alpha < 1$ : “high temperature”  
(Height of dominant paths  $h = O(n)$ )
  - ▶  $\alpha > 1$ : “low temperature”  
(Height of dominant paths  $h = O(n^0)$ )

## Phase transition

- ▶  $S_{A,\alpha}$  grows as  $O(n)$  for  $0 < \alpha < 1$  while as  $O(\ln n)$  for  $\alpha > 1$ .  
 $\Rightarrow$  Non-analytic behavior at  $\alpha = 1$  (Phase transition)
- ▶ In terms of the entanglement Hamiltonian,  
 $\text{Tr}_A \rho_A^\alpha = \text{Tr}_A e^{-\alpha H_{\text{ent}, A}}$        $\alpha$ : “inverse temperature”
  - ▶  $0 < \alpha < 1$ : “high temperature”  
 (Height of dominant paths  $h = O(n)$ )
  - ▶  $\alpha > 1$ : “low temperature”  
 (Height of dominant paths  $h = O(n^0)$ )
- ▶ The transition point  $\alpha = 1$  itself forms the third phase.





Introduction

Motzkin spin model

Colored Motzkin model

SIS Motzkin model

Colored SIS Motzkin model

Rényi entropy

Rényi entropy of Motzkin model

**Summary and discussion**

# Summary and discussion 1

## Summary

- ▶ We have reviewed the (colored) Motzkin spin models which yield large entanglement entropy proportional to the square root of the volume.

# Summary and discussion 1

## Summary

- ▶ We have reviewed the (colored) Motzkin spin models which yield large entanglement entropy proportional to the square root of the volume.
- ▶ We have extended the models by introducing additional d.o.f. based on Symmetric Inverse Semigroups.
  - ▶ Quantum phase transitions  
In uncolored case ( $\mathcal{S}_1^3$ ), log. violation v.s. area law  $O(1)$  for  $S_A$   
In colored case ( $\mathcal{S}_2^3$ ),  $\sqrt{n}$  v.s.  $\ln n$  for  $S_A$ .

# Summary and discussion 1

## Summary

- ▶ We have reviewed the (colored) Motzkin spin models which yield large entanglement entropy proportional to the square root of the volume.
- ▶ We have extended the models by introducing additional d.o.f. based on Symmetric Inverse Semigroups.
  - ▶ Quantum phase transitions
    - In uncolored case ( $\mathcal{S}_1^3$ ), log. violation v.s. area law  $O(1)$  for  $S_A$
    - In colored case ( $\mathcal{S}_2^3$ ),  $\sqrt{n}$  v.s.  $\ln n$  for  $S_A$ .
- ▶ Semigroup extension of the Fredkin model

[Padmanabhan, F.S., Korepin 2018]

# Summary and discussion 1

## Summary

- ▶ We have reviewed the (colored) Motzkin spin models which yield large entanglement entropy proportional to the square root of the volume.
- ▶ We have extended the models by introducing additional d.o.f. based on Symmetric Inverse Semigroups.
  - ▶ Quantum phase transitions
    - In uncolored case ( $\mathcal{S}_1^3$ ), log. violation v.s. area law  $O(1)$  for  $S_A$
    - In colored case ( $\mathcal{S}_2^3$ ),  $\sqrt{n}$  v.s.  $\ln n$  for  $S_A$ .
- ▶ Semigroup extension of the Fredkin model

[Padmanabhan, F.S., Korepin 2018]
- ▶ **As a feature of the extended models,** Anderson-like localization occurs in excited states corresponding to disconnected paths.
  - ▶ “2nd quantized paths”.

# Summary and discussion 2

## Summary

- ▶ We have analytically computed the Rényi entropy of half-chain in the Motzkin model.
  - ▶ Phase transition at  $\alpha = 1$  (New phase transition!)  
No analog for other spin chains investigated so far (XX, XY, AKLT,...).

# Summary and discussion 2

## Summary

- ▶ We have analytically computed the Rényi entropy of half-chain in the Motzkin model.
  - ▶ Phase transition at  $\alpha = 1$  (New phase transition!)  
No analog for other spin chains investigated so far (XX, XY, AKLT,...).
  - ▶ For  $0 < \alpha < 1$  (“high temperature”),  $S_{A,\alpha} = O(n)$ .
  - ▶ For  $\alpha > 1$  (“low temperature”),  $S_{A,\alpha} = O(\ln n)$ .

# Summary and discussion 2

## Summary

- ▶ We have analytically computed the Rényi entropy of half-chain in the Motzkin model.
  - ▶ Phase transition at  $\alpha = 1$  (New phase transition!)  
No analog for other spin chains investigated so far (XX, XY, AKLT,...).
  - ▶ For  $0 < \alpha < 1$  (“high temperature”),  $S_{A,\alpha} = O(n)$ .
  - ▶ For  $\alpha > 1$  (“low temperature”),  $S_{A,\alpha} = O(\ln n)$ .
- ▶ We also have a similar result for the Fredkin spin chain.

[F.S., Korepin, 2018]



# Summary and discussion 2

## Summary

- ▶ We have analytically computed the Rényi entropy of half-chain in the Motzkin model.
  - ▶ Phase transition at  $\alpha = 1$  (New phase transition!)  
No analog for other spin chains investigated so far (XX, XY, AKLT,...).
  - ▶ For  $0 < \alpha < 1$  (“high temperature”),  $S_{A,\alpha} = O(n)$ .
  - ▶ For  $\alpha > 1$  (“low temperature”),  $S_{A,\alpha} = O(\ln n)$ .
- ▶ We also have a similar result for the Fredkin spin chain.  
[F.S., Korepin, 2018]
- ▶ Rényi entropy of chain of general length (in progress)  
Our conjecture: the same phase transition occurs for chain of general length

# Summary and discussion 2

## Summary

- ▶ We have analytically computed the Rényi entropy of half-chain in the Motzkin model.
  - ▶ Phase transition at  $\alpha = 1$  (New phase transition!)  
No analog for other spin chains investigated so far (XX, XY, AKLT,...).
  - ▶ For  $0 < \alpha < 1$  (“high temperature”),  $S_{A,\alpha} = O(n)$ .
  - ▶ For  $\alpha > 1$  (“low temperature”),  $S_{A,\alpha} = O(\ln n)$ .
- ▶ We also have a similar result for the Fredkin spin chain. [F.S., Korepin, 2018]
- ▶ Rényi entropy of chain of general length (in progress)  
Our conjecture: the same phase transition occurs for chain of general length
- ▶ Similar computation for semigroup extensions (in progress)  
[F.S., Padmanabhan, 2018], [Padmanabhan, F.S., Korepin, 2018]

# Summary and discussion 3

## Future directions

- ▶ Continuum limit? (In particular, for colored case)

[Chen, Fradkin, Witczak-Krempa 2017]

# Summary and discussion 3

## Future directions

- ▶ Continuum limit? (In particular, for colored case)

[Chen, Fradkin, Witczak-Krempa 2017]

- ▶ Holography? Application to quantum gravity or black holes?

[Alexander, Klich 2018]

- ▶ Higher-dimensional models ( $d = 2, 3, \dots$ )?

# Summary and discussion 3

## Future directions

- ▶ Continuum limit? (In particular, for colored case)

[Chen, Fradkin, Witczak-Krempa 2017]

- ▶ Holography? Application to quantum gravity or black holes?



[Alexander, Klich 2018]


- ▶ Higher-dimensional models ( $d = 2, 3, \dots$ )?

Thank you very much for your attention!

- ▶ By adding the balancing term to the Hamiltonian

$$\lambda_2 \sum_{j=1}^{2n-1} \left[ \prod |(x_{1,3})_j, (x_{3,2})_{j+1}\rangle + \prod |(x_{2,3})_j, (x_{3,1})_{j+1}\rangle \right]$$

with  $\lambda_1$  put to the term   $\sim$  , quantum phase transition takes place in the 4 sectors except (3, 3):



$\lambda_1, \lambda_2 > 0$  is not frustration free (here, we do not consider).