

CHARACTERIZING THE INCREASE OF THE RESIDUAL ORDER UNDER BLOWUP IN POSITIVE CHARACTERISTIC

HERWIG HAUSER, STEFAN PERLEGA

ABSTRACT. In contrast to the characteristic zero situation, the residual order of an ideal may increase in positive characteristic under permissible blowups at points of the exceptional divisor where the order of the ideal has remained constant. The specific situations where this happens are described explicitly.

1. INTRODUCTION

To prove embedded resolution of singularities in characteristic zero for a reduced subscheme X of a regular ambient scheme W equipped with a normal crossings divisor D one typically associates to every point a of X a local invariant $\text{inv}_a X$ measuring the complexity of the singularity of X at a and the position of X with respect to D . The invariant consists of a string of non-negative integers, is upper semicontinuous and decreases lexicographically when X is blown up along the center Z defined as the locus of points where $\text{inv}_a X$ attains its maximal value. This is done in a way so that Z is regular and has normal crossings with D . As the invariant varies in the well ordered set $(\mathbb{N}^N, \text{lex})$ and its minimal value corresponds to a regular point a at which X has normal crossings with D , the resolution of X is obtained by induction [Hir64, Vil89, Vil92, BM97, EH02, Cut04, Wł05, Kol07].

For the first component of $\text{inv}_a X$ the simplest choice is the order $\text{ord}_a J$ of the defining ideal J of X in W . Blowing up a regular center contained in the associated equimultiple locus of J , the order does not increase, $\text{ord}_{a'} J' \leq \text{ord}_a J$, for all points a' in the weak transform X' of X above a . At a point a' where the order remains constant, the second component of $\text{inv}_a X$ comes into play. Leaving aside transversality issues of Z with D , it is (usually) defined as the order of the coefficient ideal K of J at a with respect to a hypersurface of maximal contact V , less the exceptional multiplicity of K . This numeral, which we will call the residual order of J at a (the name *residual order* was introduced by Hironaka for the situation in positive characteristic in [Hir12]), does not depend on the choice of the hypersurface and is again upper semicontinuous along the strata defined by the order of J . It is thus well suited to form the second component of $\text{inv}_a X$. Blowing up a regular center Z inside the top loci of the order and the residual order, the second component does not increase whenever the first remains constant. From this point on, the argument is repeated until, by exhaustion of dimensions, a decrease of the invariant under blowup is established.

This approach to resolution has several drawbacks in positive characteristic: First, hypersurfaces of maximal contact no longer exist; a possible substitute are

hypersurfaces of *weak* maximal contact as introduced in [EH02, Hau03, Hau10]. These are defined as regular hypersurfaces maximizing the order of the coefficient ideal (in characteristic zero, the maximum can be realized by a hypersurface of maximal contact.) Secondly, the resulting residual order is no longer upper semi-continuous, so its top locus need not be closed; extra care has to be taken. Finally, even if centers are chosen appropriately, the residual order may still go up under blowup at points where the order of the ideal J has remained constant. This increase is also known as the “kangaroo phenomenon”. It destroys the induction argument.

In view of these difficulties, two approaches are plausible: Either to reject the residual order as a valuable resolution invariant in positive characteristic and to search for new invariants. This option has been undertaken with a certain success by several authors [Hir84, Cos87, Cos91, Vil07, Kaw07, KM10]. Or, to try to understand better the circumstances where the residual order behaves badly in order to develop an exit strategy for the obstructions. This is the proposal we wish to pursue in the present paper.

In this spirit, the situations where an increase of the residual order occurs under blowup with permissible choices of centers will be investigated in detail. It turns out that in order to produce an increase, the defining equations of X in W must satisfy quite restrictive conditions: The (weighted) initial forms of minimal order of the elements of J have a unique form (up to constant factors and coordinate changes), and are actually powers of purely inseparable polynomials. Their logarithmic Hasse derivatives have a specific shape, and the exceptional multiplicities of the coefficient ideal K of J satisfy an explicit arithmetic inequality. These three conditions are satisfied simultaneously only in very special cases.

Along the proof of these facts, we extend Moh’s bound on the possible increase of the residual order to non-hypersurfaces and not necessarily purely inseparable power series [Moh87]. The upshot of the results is as follows (see section 3 for the precise statements):

Theorem. *Let J be an ideal of W of order c at a , with coefficient ideal K of order o with respect to a hypersurface of weak maximal contact. Assume that the residual order of J with respect to a given normal crossings divisor D increases under permissible blowup at a point a' where c has remained constant. Then c is a multiple $m \cdot p^e$ of a power of the characteristic p , o is a multiple $w \cdot c!$ of $c!$, the weighted initial form with respect to w of elements f of J of minimal weighted order is a power $\text{in}_w(f) = (z^c + F(x))^m$, with F a homogeneous polynomial of degree $w \cdot p^e$ in variables x_1, \dots, x_n , not a p^e -th power, and with a specific logarithmic Hasse derivative $x_k^{p^\ell} \partial_{x_k} F$. Here, $\ell < e$ is maximal so that F is a p^ℓ -th power.*

Choosing x_i subordinate to D and factorizing F maximally into $F = x^r \cdot G$, the residues modulo $p^{\ell+1}$ of the exceptional exponents r_i satisfy $\sum \bar{r}_i \leq (b-1) \cdot p^{\ell+1}$, where the sum ranges over those exceptional components which are lost when passing from a to a' , and where b is the number of $r_i \not\equiv p^{\ell+1}$ among them.

In the above situation, the residual order of J increases at most by $\frac{c!}{p}$.

Various other notable approaches to the resolution problem in positive characteristic can be found in [Abh56, Gir75, Hir84, Cos87, Cos91, Moh96, Cut04, Cut11, Hau04, HW14, Vil07, Kaw07, KM10].

Acknowledgments. We are indebted to the following mathematicians for many valuable discussions and suggestions: S. Abhyankar, H. Hironaka, G. Müller, J. Schicho, A. Quirós, S. Encinas, O. Villamayor, A. Bravo, D. Cutkosky, J.-J. Risler, V. Cossart, J. Włodarczyk, H. Kawanoue, K. Matsuki, D. Panazzolo, M. Spivakovsky, F. Cano, R. Blanco, D. Zeillinger, D. Wagner, A. Frühbis-Krüger.

2. SETTING

The concepts and constructions that are successfully used to prove the embedded resolution of singularities over fields of characteristic zero require some amendments for their characteristic free definition. It remains an open problem whether these will suffice to give a proof of resolution in positive characteristic and arbitrary dimension.

We shall work with complete regular local rings $R = (R, m_R)$ of dimension $n + 1$ over an algebraically closed field \mathbb{K} , and of residue field $R/m_R \cong \mathbb{K}$. By Cohen's structure theorem, R is a formal power series ring in $n + 1$ variables over \mathbb{K} . It should be thought of as the completion of the local ring of some regular noetherian scheme W over \mathbb{K} at a closed point a , and ideals J of R as defining the formal neighborhood at a of a closed subscheme X of W . Typically, a regular system of parameters $(z, x) = (z, x_1, \dots, x_n)$ will be chosen, with a distinguished parameter z . We then often fix a ring inclusion $\rho : R/(z) \cong Q \subset R$ providing a section of the projection $R \rightarrow R/(z)$, for some subring Q of R . The induced isomorphism $R \cong Q[[z]]$ will be used frequently.

The *order* of an ideal J in R is defined as $\text{ord } J = \text{ord}_{m_R} J = \sup\{k \in \mathbb{N}, J \subset m_R^k\}$. If P is a prime ideal of R , we define the order $\text{ord}_P J$ of J with respect to P as the order of $J \cdot R_P$ in the localization R_P . For regular ideals P , it equals $\sup\{k \in \mathbb{N}, J \subset P^k\}$. A closed subscheme $Z = V(P)$ of $\text{Spec}(R)$ is said to be *contained in the equimultiple locus of J* if $\text{ord}_P J = \text{ord } J$ holds.

The *initial form* $\text{in}(f)$ of an element $f \in R$ is the homogeneous polynomial of lowest degree of the power series expansion of f with respect to the m_R -adic filtration of R . If z, x_1, \dots, x_n are given regular parameters in R and $w \in \mathbb{Q}_{>0}$ is a positive rational number, we define for $f \in R$ with expansion $f(z, x) = \sum_{i \geq 0} f_i z^i$ and coefficients $f_i \in \mathbb{K}[[x_1, \dots, x_n]]$ the *weighted order* $\text{ord}_w f$ of f with respect to the weight vector $(w, 1, \dots, 1)$ as the minimum of the values $wi + \text{ord } f_i$, the order of f_i being taken in $\mathbb{K}[[x_1, \dots, x_n]]$. It clearly only depends on the choice of z . The *weighted initial form* $\text{in}_w(f)$ of f with respect to the weight vector $(w, 1, \dots, 1)$ (and the parameters (z, x)) is then defined as the sum $\text{in}_w(f) = \sum \text{in}(f_i) z^i$, where the sum ranges over those i for which the minimal value of $wi + \text{ord } f_i$ is attained.

Let $\pi : R \rightarrow R'$ be a *completed local blowup* of R , with regular center $Z = V(P)$ in $W = \text{Spec}(R)$, for some ideal P of R and a complete regular local ring R' . By this we understand that R' is the completion of a local ring $\mathcal{O}_{W', a'}$ where $W' = \text{Proj}(\oplus_{i \geq 0} P^i)$ is the blowup of W in Z , $a' \in W'$ is a closed point and $\pi : R \rightarrow R'$ is the induced map of complete local rings. As R and Z are regular, R' is again regular, and actually isomorphic to R . Occasionally we shall identify R' with R .

The *weak transform* of an ideal J of R under π is defined as the (unique) ideal J' of R' so that

$$\pi(J) \cdot R' = x_1^{\text{ord}_P J} \cdot J',$$

where $x_1 \in R'$ defines the exceptional component E of π in $\text{Spec}(R')$. It is well known that under blowups in regular centers contained in the equimultiple locus of J the order of J does not increase when passing to J' [Hau14].

Define the *coefficient ideal* $K = \text{coeff}_V(J)$ of J with respect to a regular hypersurface $V = V(z)$ in $\text{Spec}(R)$ and a section $\rho : R/(z) \cong Q \subset R$ of $R \rightarrow R/(z)$ as the ideal of Q defined by

$$\text{coeff}_V(J) = \sum_{i < c} (f_i, f \in J)^{\frac{c-i}{c}},$$

where $c = \text{ord } J$ and elements $f \in J$ are expanded as series $f = \sum_{i \geq 0} f_i z^i$ in $Q[[z]]$, with coefficients f_i in Q . The coefficient ideal depends on V and ρ , but not on the choice of the parameter z defining V . By abuse of notation, we suppress the dependence of the coefficient ideal on the choice of the section ρ . This does no harm in our context since the order of the coefficient ideal (which is our main concern) only depends on V and not on ρ .

Let $V = V(z)$ be a regular hypersurface in $\text{Spec}(R)$ and let o be the order of the coefficient ideal $\text{coeff}_V(J)$. Further, set $w = \frac{o}{c}$ where c is the order of the ideal J . Then the minimum of the weighted orders $\text{ord}_w(f)$ of elements $f \in J$ with respect to the regular parameter z and the weight vector $(w, 1, \dots, 1)$ equals $c \cdot w$.

A regular hypersurface $V = V(z)$ in $\text{Spec}(R)$ has *weak maximal contact with J* if the order of the coefficient ideal $\text{coeff}_V(J)$ of J with respect to V is maximized over all choices of regular hypersurfaces in $\text{Spec}(R)$ and if for any blowup with regular center $Z = V(P)$ contained in V and in the equimultiple locus of J , the strict transform of V in $\text{Proj}(\oplus_{i \geq 0} P^i)$ contains all points at which the order of the weak transform of J has remained constant.

Two cases can occur: The supremum of the orders of $\text{coeff}_V(J)$ over all V may be infinite, in which case J is of the form $J = (z^c)$ for some regular parameter $z \in R$, and has trivial coefficient ideal equal to 0 with respect to $V = V(z)$. This case is irrelevant for our investigations and will be discarded. Or, the supremum of the orders is bounded, in which case the maximum exists and is realized by some V . Such a V can then be chosen so that its strict transform contains all points where the order of the weak transform of J has remained constant. If the characteristic is zero, then V can even be chosen in a way so that it has maximal contact with J .

Let D be a (not necessarily reduced) normal crossings divisor in $\text{Spec}(R)$, and let $J \subset R$ be an ideal. A regular hypersurface $V = V(z)$ is *compatible* with D and J if it has normal crossings with D and if there is a section $\rho : R/(z) \cong Q \subset R$ of $R \rightarrow R/(z)$ so that the coefficient ideal $K = \text{coeff}_V(J)$ of J with respect to V and ρ factors into $K = M \cdot I$, for some ideal I of Q , where M is the principal ideal of Q defining $D \cap V$ in V .

Let J be an ideal and D a normal crossings divisor for which there exists a regular hypersurface V that has weak maximal contact with J and is compatible with D . The *residual order* of such an ideal J with respect to D is defined as

$$\text{residual-order}_D(J) = \text{ord}(\text{coeff}_V(J)) - \text{ord } M = \text{ord } I,$$

where $V = V(z) \subset \text{Spec}(R)$ is a hypersurface of weak maximal contact with J and compatible with D , and where M is the ideal which defines $D \cap V$ in V and appears in the factorization $\text{coeff}_V(J) = M \cdot I$. Notice that the residual order is independent of the choice of V . This numeral is frequently used in the proof of resolution of singularities in characteristic zero. It is supposed to measure the “distance” of K from being a principal monomial ideal supported by D .

A completed local blowup $\pi : R \rightarrow R'$ with center $Z = V(P)$ is said to be *permissible* with respect to J and D if the center Z of π is regular, has normal crossings with D and if there exists a hypersurface V of weak maximal contact with J , compatible with D , and such that Z is contained in V and in the equimultiple loci of J and I ; here I is defined through $\text{coeff}_V(J) = M \cdot I$ as before.

The *transform* D' of D with respect to J under a permissible completed local blowup $\pi : R \rightarrow R'$ is defined as the normal crossings divisor $D' = D^s + (\text{ord}_P K - c!) \cdot E$ in $\text{Spec}(R')$, where D^s denotes the strict transform of D and $E = \pi^{-1}(Z)$ is the new exceptional component (cf. [EH02]). Here, c is the order of J in R , and K is the coefficient ideal of J with respect to a hypersurface of weak maximal contact V with J and compatible with D . The definition of D' is independent of the choice of the hypersurface V of weak maximal contact.

If J and D admit a regular hypersurface V having weak maximal contact with J and compatible with D , it can be shown that there exists, for every permissible completed local blowup $\pi : R \rightarrow R'$ under which the order of J remains constant, a regular hypersurface U' in $\text{Spec}(R')$ which has weak maximal contact with the weak transform J' of J and is compatible with D' (cf. the proof of the proposition below). If the characteristic is zero, then the hypersurface V in $\text{Spec}(R)$ can be chosen in such a way that its strict transform V' in $\text{Spec}(R')$ has these properties. This is no longer true over fields of positive characteristic.

Regular parameters (z, x) in R are called *subordinate* to a permissible blowup π , an ideal J , a normal crossings divisor D and a hypersurface V of weak maximal contact with J and compatible with D , if $V = V(z)$, the components of D are supported by the hypersurfaces $V(x_i)$ of $\text{Spec}(R)$, and if the defining ideal P of the center Z is generated by z and x_i , for i varying in a subset S of $\{1, \dots, n\}$. Permuting the x_i if necessary, we may assume that the blowup occurs in the x_1 -chart. There then exist a subset T of S containing 1 and constants $t_i \in \mathbb{K}^*$, for $i \in T \setminus \{1\}$, so that π is defined by

$$\begin{aligned} z &\rightarrow x_1 z, \\ x_1 &\rightarrow x_1, \\ x_i &\rightarrow x_1(x_i + t_i) \quad \text{for } i \in T \setminus \{1\}, \\ x_i &\rightarrow x_1 x_i \quad \text{for } i \in S \setminus T, \\ x_i &\rightarrow x_i \quad \text{for } i \notin S, \end{aligned}$$

where R' is identified with R and (z, x_1, \dots, x_n) denote regular parameter systems in R and R' . We may further assume that either $T = \{1\}$ or that for all indices $i \in T$ the inclusion $V(x_i) \subset D$ holds.

If the characteristic of \mathbb{K} is zero, it is well-known that for all permissible completed local blowups $\pi : R \rightarrow R'$ under which the order of J remains constant $\text{ord } J' = \text{ord } J$ when passing to its weak transform J' , the residual order does not increase, i.e.,

$$\text{residual-order}_{D'} J' \leq \text{residual-order}_D J$$

holds. Over fields of positive characteristic, this is no longer true: the residual order may increase.

3. RESULTS

The characterization of ideals and permissible blowups for which the residual order increases goes as follows.

Theorem. *Let R be a complete regular local noetherian ring R of dimension $n + 1$ over an algebraically closed field \mathbb{K} of positive characteristic $p > 0$. Let D be a normal crossings divisor in $\text{Spec}(R)$. Let be given an ideal J in R admitting a hypersurface of weak maximal contact and compatible with D . Let $\pi : R \rightarrow R'$ be a completed local blowup of R , permissible with respect to J and D with center Z defined by the ideal P of R . Denote by J' the weak transform of J in R' , and by D' the transform of D in $\text{Spec}(R')$ with respect to J .*

Let V in $\text{Spec}(R)$ be a hypersurface of weak maximal contact with J and compatible with D such that Z is contained in V and in the equimultiple loci of J and I , where I appears in the factorization $\text{coeff}_V(J) = M \cdot I$, with M the ideal in V defining $D \cap V$. Choose regular parameters $(z, x) = (z, x_1, \dots, x_n)$ of R subordinate to π , J , D and V , and let $T \subset S \subset \{1, \dots, n\}$ and $t_i \in \mathbb{K}^$ be as above.*

Assume that J' has the same order as J but that its residual order with respect to D' is larger than the residual order of J with respect to D . Then the following conditions must be satisfied.

- (1) *The order c of J is a multiple $c = m \cdot p^e$ of a p -th power, with $m \geq 1$ not divisible by p and $e \geq 1$.*
- (2) *The order o of the coefficient ideal K of J with respect to V is a multiple $o = w \cdot c!$ of $c!$, with $w \geq 2$.*
- (3) *There exists a homogeneous, non p^e -th power polynomial F in x_1, \dots, x_n of degree $w \cdot p^e$ so that the weighted initial form $\text{in}_w(f)$ with respect to (z, x) and the weights $(w, 1, \dots, 1)$ of every element $f \in J$ of minimal weighted order wc is the m -th power of a purely inseparable polynomial, say*

$$\text{in}_w(f) = \alpha \cdot (z^{p^e} + F(x))^m,$$

for some non-zero constant $\alpha \in \mathbb{K}^$.*

- (4) *Factorize F into $F(x) = x^r \cdot G(x)$ with $r_i = \text{ord}_{(x_i)} F$, for $i \in T$, and G a homogeneous polynomial of degree $v = \deg F - \sum_{i \in T} r_i$. If Q_T denotes the ideal of $\mathbb{K}[[x_1, \dots, x_n]]$ generated by $x_i - t_i x_1$, for $i \in T \setminus \{1\}$, and x_i , for $i \notin T$, then*

$$\text{ord}_{Q_T}^{\text{mod } p^e} F > \deg G,$$

where $\text{ord}_{Q_T}^{\text{mod } p^e} F$ denotes the maximum of the orders $\text{ord}_{Q_T}(F + H^{p^e})$ over all polynomials H in x_1, \dots, x_n .

The inequality $\text{ord}_{Q_T}^{\text{mod } p^e} F > \deg G$ from (4) implies the following conditions (5) to (9). Let $\ell < e$ be the largest integer so that F is a p^ℓ -th power, and denote by b the number of exponents r_i , for $i \in T$, not congruent to $p^{\ell+1}$.

- (5) *Denote by \mathfrak{t} the vector in \mathbb{K}^n of components t_i for $i \in T \setminus \{1\}$, and 0 otherwise. The polynomial $G(x)$ of the factorization $F(x) = x^r \cdot G(x)$ has, up to p^e -th powers, a unique form,*

$$G((1, x_2, \dots, x_n) + \mathfrak{t}) = \lfloor \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i} \cdot N^{p^e}(x_2, \dots, x_n) \rfloor_v$$

for some polynomial $N(x_2, \dots, x_n)$. Here, the product $\prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i}$ is considered as a power series, and $\lfloor - \rfloor_v$ denotes the v -jet of a power series.

- (6) The residues $0 \leq \bar{r}_i < p^{\ell+1}$ of r_i modulo $p^{\ell+1}$ satisfy the arithmetic inequality

$$\sum_{i \in T} \bar{r}_i \leq (b-1) \cdot p^{\ell+1}.$$

Equivalently, one has

$$\sum_{i \in T} \bar{r}_i + \bar{u} \neq b \cdot p^{\ell+1}.$$

- (7) For $j \notin T$, the variables x_j appear only as p^e -th powers in $F(x)$, say

$$F(x) \in \mathbb{K}[x_i^{p^\ell}, x_j^{p^e}, i \in T, j \notin T].$$

- (8) For $i \in T$, the p^ℓ -th logarithmic Hasse derivatives of $F(x)$ with respect to x_i are of the form

$$x_i^{p^\ell} \cdot \partial_{x_i^{p^\ell}} F(x) = x^r \cdot H_i(x),$$

where H_i is a polynomial in $(x_j - t_j x_1)^{p^\ell}$ and $x_k^{p^e}$, for $j \in T \setminus \{1\}$ and $k \notin T$.

- (9) The increase of the residual order is bounded by

$$\text{residual-order}_{D'} J' \leq \text{residual-order}_D J + \frac{c!}{p}.$$

Comments. The statements of the theorem crystallize a broader background which will be explained below.

(a) The theorem only tells us something about the exceptional multiplicities and the (weighted) tangent cone of the ideal J . It does not make any statement about the higher order terms of the elements of J .

(b) The multiplicity of the new exceptional component in D' equals $\text{ord}_P K - c!$ and is hence a multiple of $c!$. Let x_1 be the parameter defining this component. Then the center $Z' = V(z, x_1)$ in $\text{Spec}(R')$ is contained in the equimultiple locus of J' , has normal crossings with D' and can be blown up until the exceptional multiplicity of this component has dropped to 0.

(c) The residual order is a questionable resolution invariant as is exhibited by Perlega's example of an infinite sequence of permissible blowups where the residual order tends to infinity [HP16b]. In this sequence, however, the centers are not chosen of maximal dimension, so this is not yet a counterexample to the resolution of singularities in positive characteristic.

(d) The increase of the residual order can only happen if under the blowup at least two components of D are lost when passing to the reference point a' in the new exceptional component.

(e) The increase of the residual order represents a serious obstacle for trying to transfer the proof of resolution of singularities in characteristic zero to positive characteristic. For surfaces, it can still be used, but has to be modified slightly so as to perform appropriately under blowup, see [HW14, HP16a]. Already for threefolds the situation is unclear and no efficient resolution invariant (for embedded resolution) seems to be known (for the non-embedded case, see [Abh66, CP08, CP09, Cut09]).

(f) For a fixed prime number p , the arithmetic inequality for the residues of the multiplicities r_i in assertion (6) of the theorem always holds when T contains sufficiently many indices i with $r_i \not\equiv p^{\ell+1}$.

(g) For fixed numbers n, p, e and ℓ as in the theorem, a homogeneous polynomial $F(x) = x^r \cdot G(x)$ of degree divisible by p^e , but not a p^e -th power, defines via $f = z^{p^e} + F(x)$ a weighted homogeneous hypersurface singularity whose residual order increases under blowup if and only if $G(x)$ is of the form specified in assertion (5) and the multiplicities r_i fulfill the arithmetic inequality in assertion (6) of the theorem.

(h) Assertion (2) and the bound in (9) have been known to Moh in the case of a purely inseparable hypersurface singularity [Moh87].

4. AUXILIARY RESULTS

The proof of the theorem will rely on the following more technical result.

Proposition. *In the situation of the theorem, there exists an automorphism of R sending z onto $u = z - q$, for some $q \in Q$ of order $\text{ord } q \geq \frac{o}{c_1} > 1$, and inducing the identity on Q , so that $U' = V(u')$ has again weak maximal contact with J (but may no longer be compatible with D), and so that the following two conditions are satisfied.*

- (1) *The strict transform U' of U has weak maximal contact with J' and is compatible with D' .*
- (2) *Factorize the coefficient ideal $K'_1 = \text{coeff}_{U'}(J')$ of J' with respect to U' and $\sigma'_1 : R'/(u') \cong Q \subset R'$ into $K'_1 = M'_1 \cdot I'_1$ with M'_1 the principal monomial ideal defining $D' \cap U'$ in U' . Then the residual order $\text{ord } I'_1$ of J' with respect to D' is bounded by*

$$\text{ord } I'_1 \leq \text{ord}_{Q_T} \text{in}(h) - \sum_{i \notin T} s_i,$$

for any element h of minimal order o of the coefficient ideal $K_1 = \text{coeff}_U(J)$ of J with respect to U and the ring inclusion $\sigma_1 : R/(u) \cong Q \subset R$, and where $s_i = \text{ord}_{(x_i)} D$.

To show this, we need two lemmata. Lemma 1 will clarify how the orders of the coefficients f_i in the expansion $f = \sum_{i \geq 0} f_i z^i$ are related to the order of the coefficient ideal. In Lemma 2 we will investigate the effect of coordinate changes $u = z - q$ with $q \in \mathbb{K}[[x]]$ on the coefficient ideal. In particular, we will see that if the coordinate change increases the order of the coefficient ideal with respect to $V(z)$, then the element q has to be of a very specific form.

Lemma 1. *Let R be the power series ring $\mathbb{K}[[z, x]]$ with $x = (x_1, \dots, x_n)$ for some field \mathbb{K} . Denote the maximal ideal of R by m_R . Let $J \subset R$ be an ideal of order $\text{ord } J = c$. Let each element $f \in J$ have the expansion $f = \sum_{i \geq 0} f_i z^i$ with $f_i \in \mathbb{K}[[x]]$.*

Set $K = \text{coeff}_V(J)$ for $V = V(z)$ and a section $\rho : R/(z) \cong Q \subset R$ of $R \rightarrow R/(z)$. Define $o = \text{ord } K$ and $w = \frac{o}{c_1}$.

Then the following statements hold:

- (1) *The order of K can be expressed as*

$$o = \min_{f \in J} \min_{i < c} \frac{c!}{c-i} \text{ord } f_i.$$

Consequently, for all elements $f \in J$ and indices $i < c$, the inequality

$$\text{ord } f_i \geq (c - i)w$$

holds.

$$(2) \quad o \geq c!$$

$$(3) \quad o > c! \text{ holds if and only if } J \equiv (z^c) \text{ modulo } m_R^{c+1}.$$

Proof. Immediate from the definition of the coefficient ideal. \square

Lemma 2. Let R be the power series ring $\mathbb{K}[[z, x]]$ with $x = (x_1, \dots, x_n)$ where \mathbb{K} is a field of characteristic $p > 0$. Consider a change of coordinates $u = z - q$ where $q \in \mathbb{K}[x]$ is a homogeneous polynomial, and define $V = V(z)$, $U = V(u)$. Let $J \subset R$ be any ideal of order $\text{ord } J = c$ and let p^e the largest p -th power dividing c .

Let $K = \text{coeff}_V(J)$ and $K_1 = \text{coeff}_{V_1}(J)$, and set $o = \text{ord } K$, $o_1 = \text{ord } K_1$, and $w = \frac{o}{c!}$. The following statements hold:

$$(1) \quad \text{If } \deg q \geq w, \text{ then } o_1 \geq o.$$

$$(2) \quad \text{If } \deg q < w \text{ and there exists an element } f \in J \text{ that is } z\text{-regular of order } c, \text{ then } o_1 = c! \cdot \deg q < o.$$

$$(3) \quad \text{Let } 1 \leq i \leq n \text{ be an index. If } \text{ord}_{(x_i)} q \geq \frac{1}{c!} \text{ord}_{(x_i)} K, \text{ then } \text{ord}_{(x_i)} K_1 \geq \text{ord}_{(x_i)} K.$$

$$(4) \quad \text{Let } f \in J \text{ be an element that is } z\text{-regular of order } c. \text{ Let } f \text{ have the expansion } f = \sum_{i \geq 0} f_i z^i \text{ with } f_i \in \mathbb{K}[[x]]. \text{ If } o_1 > o \text{ holds, then } \deg q = w \text{ and } q \text{ fulfills}$$

$$q^{p^e} = \lambda \cdot \text{in}(f_{c-p^e})$$

for a non-zero constant $\lambda \in K^*$.

Proof. Let each element $f \in J$ have expansions $f = \sum_{i \geq 0} f_i z^i$ and $f = \sum_{i \geq 0} \tilde{f}_i u^i$ with $f_i, \tilde{f}_i \in \mathbb{K}[[x]]$. Then

$$\tilde{f}_i = \sum_{k \geq i} \binom{k}{i} f_k q^{k-i}.$$

Notice that an element f is z -regular of order c if and only if the coefficient f_c is a unit.

Statements (1), (2) and (3) can be verified directly by using the formula for \tilde{f}_i and Lemma 1 (1).

To prove statement (4), let $f \in J$ be z -regular of order c . If $\deg q > w$, it is straightforward to show that $o_1 = o$ holds. By statement (2) this implies that $\deg q = w$ has to hold.

Assume now that there exists an index $c - p^e < i < c$ such that $\text{ord } f_i = (c - i)w$. Let i be maximal with this property. Notice that $\binom{c}{i} \equiv 0 \pmod{p}$ by Lucas' theorem on binomial coefficients in characteristic $p > 0$. Using the form of \tilde{f}_i , the maximality of i and the fact that $\binom{c}{i} \equiv 0 \pmod{p}$, it follows that $\text{ord } \tilde{f}_i = (c - i)w$. Consequently, $o_1 \leq o$ holds by Lemma 1 (1), contradicting the assumption.

Hence, $\text{ord } f_i > (c - i)w$ holds for all indices $c - p^e < i < c$. If we had $\text{ord } f_{c-p^e} > p^e w$, then

$$\text{in}(\tilde{f}_{c-p^e}) = \binom{c}{p^e} f_c(0) q^{p^e}.$$

Since $\binom{c}{p^e} \not\equiv 0 \pmod{p}$ by Lucas' theorem, this implies that $\text{ord } \tilde{f}_{c-p^e} = p^e w$ and hence, $o_1 \leq o$ by Lemma 1 (1), again a contradiction.

Thus, $\text{ord } f_{c-p^e} = p^e w$. Since $o_1 > o$, it follows that $\text{ord } \tilde{f}_{c-p^e} > p^e w$. This gives

$$\text{in}(f_{c-p^e}) + \binom{c}{p^e} f_c(0) q^{p^e} = 0,$$

which proves the assertion. \square

Proof of the proposition: Define parameters $y = (y_1, \dots, y_n)$ by setting

$$y_i = \begin{cases} x_i - t_i x_1 & \text{for } i \in T \setminus \{1\}, \\ x_i & \text{otherwise.} \end{cases}$$

Notice that the ideal P is generated by the parameters z and y_i for $i \in S$. Also, $Q_T = (y_2, \dots, y_n)$. Further, the map $\pi : R \rightarrow R'$ has the following simple form with respect to (z, y) : $z \rightarrow x_1 z$, $y_1 \rightarrow x_1$, $y_i \rightarrow x_1 x_i$ for $i \in S \setminus \{1\}$ and $y_i \rightarrow x_i$ for $i \notin S$. One also says that the blowup map $\pi : R \rightarrow R'$ is monomial with respect to the parameters (z, y) . Monomial blowup maps have the advantage that they make calculations in coordinates particularly easy.

We will begin with assertion (1). To this end, let us first verify that $J \equiv (z^c)$ modulo m_R^{c+1} .

Recall that the strict transform $V' = V(z)$ of V in $\text{Spec}(R')$ is non-empty since $\text{ord } J' = \text{ord } J$ holds and V has weak maximal contact with J . Further, it is easy to see that the regular hypersurface V' is compatible with D' .

Recall that the center of blowup Z is contained in the equimultiple locus of I . Thus, it is easy to see that the ideal I' in the factorization $\text{coeff}_{V'}(J') = M' \cdot I'$ fulfills $\text{ord } I' \leq \text{ord } I$. Since we assumed that the residual order increases under the local blowup π , we conclude from this that V' does not have weak maximal contact with J' .

Assume now that $J \not\equiv (z^c)$ modulo m_R^{c+1} . Since V has weak maximal contact with J , this implies by Lemma 1 (3) that $\text{ord } \text{coeff}_{\tilde{V}}(J) = c!$ holds for all regular hypersurfaces $\tilde{V} \subset \text{Spec}(R)$. Consequently, by Lemma 1 (3)

$$\dim_{\mathbb{K}} \left(J + m_R^{c+1} / m_R^{c+1} \right) \geq 2.$$

It is straightforward to verify that this implies

$$\dim_{\mathbb{K}} \left(J' + m_{R'}^{c+1} / m_{R'}^{c+1} \right) \geq 2.$$

This would imply by Lemma 1 that any regular hypersurface $\tilde{U} \subset \text{Spec}(R')$ has weak maximal contact with J' . Since we already know that V' does not have weak maximal contact with J' , this is a contradiction. Hence, $J \equiv (z^c)$ modulo m_R^{c+1} holds.

It is immediate to see that this implies the existence of an element $f \in J$ which is z -regular of order c . Set $f' = x_1^{-c} \pi(f) \in J'$. Then the element f' is also z -regular of order c . Let these elements have the expansions $f = \sum_{i \geq 0} f_i z^i$ and $f' = \sum_{i \geq 0} f'_i z^i$ with $f_i, f'_i \in \mathbb{K}[[x]]$.

Let $\tilde{U} = V(\tilde{u}) \subset \text{Spec}(R')$ be a regular hypersurface which has weak maximal contact with J' . Set $\tilde{K} = \text{coeff}_{\tilde{U}}(J')$. If the element $\tilde{u} \in R'$ is not z -regular, then it is straightforward to verify with Lemma 1 (1) and using the fact that f is z -regular of order c that $\text{ord } \tilde{K} = c!$. This contradicts the fact that V' does not have weak maximal contact with J' . Thus, we may assume that $\tilde{u} = z + \tilde{g}$ for an element $\tilde{g} \in \mathbb{K}[[x]]$.

Set $K' = \text{coeff}_{V'}(J')$. By Lemma 2 we know that $\text{ord } \tilde{g} = \frac{1}{c!} \text{ord } K'$. Set $g = \text{in}(\tilde{g})$. Further, define $u' = z - g$, $U' = V(u')$ and $K'_1 = \text{coeff}_{U'}(J')$. Then it is clear by Lemma 2 that the chain of inequalities

$$\text{ord } \tilde{K} \geq \text{ord } K'_1 > \text{ord } K'$$

holds. We know by Lemma 2 (4) that

$$g^{p^e} = \lambda \cdot \text{in}(f'_{c-p^e})$$

for some non-zero constant λ , where p^e is the largest p -th power dividing c . Notice that $f'_{c-p^e} = x_1^{-p^e} \pi(f_{c-p^e})$.

It is clear that for all indices $i = 1, \dots, n$ the inequality

$$\text{ord}_{(x_i)} g \geq \frac{1}{p^e} \text{ord}_{(x_i)} f'_{c-p^e} \geq \frac{1}{c!} \text{ord}_{(x_i)} K'$$

holds. Since $V' = V(z)$ is compatible with D' , this implies by Lemma 2 (3) that U' is also compatible with D' .

Since the map $\pi : R \rightarrow R'$ is monomial with respect to the parameters (z, y) , it is straightforward to verify that there exists an element $q \in R$ that fulfills $g = x_1^{-1} \pi(q)$ and is of the form

$$q^{p^e} = \lambda \cdot \text{in}_\sigma(f_{c-p^e})$$

where $\text{in}_\sigma(f_{c-p^e})$ denotes the weighted initial form of f_{c-p^e} with respect to the weights $\sigma(y_i) = 2$ for $i \in S \setminus \{1\}$ and $\sigma(y_i) = 1$ for all other indices i . Set $u = z - q$. Then

$$\pi(u) = x_1(z - q) = x_1 u'.$$

Hence, the hypersurface U' is the strict transform of the hypersurface $U = V(u)$. Set $K_1 = \text{coeff}_U(J)$.

We will now show that U has weak maximal contact with J and K_1 has a factorization

$$K_1 = \left(\prod_{i \notin T} x_i^{s_i} \right) \cdot I_1$$

for some ideal I_1 of Q . Since $q^{p^e} = \text{in}_\sigma(f_{c-p^e})$, it is clear that the inequality

$$\text{ord } q \geq \frac{1}{p^e} \text{ord } f_{c-p^e} \geq \frac{o}{c!}$$

holds, as well as, for all indices $i \notin T$, the inequalities

$$\text{ord}_{(x_i)} q \geq \frac{1}{p^e} \text{ord}_{(x_i)} f_{c-p^e} \geq \frac{1}{c!} \text{ord}_{(x_i)} K$$

(since $x_i = y_i$ for $i \notin T$). By Lemma 2 (1) and (3) this implies that

$$\text{ord } K_1 \geq \text{ord } K = o$$

holds and that K_1 has the claimed factorization. Since the hypersurface V already had weak maximal contact with J , we must have $\text{ord } K_1 = o$. Hence, also U has weak maximal contact with J .

We now consider two cases. If the hypersurface U' has weak maximal contact with J' , we are done. On the other hand, if $\text{ord } \tilde{K} > \text{ord } K'_1$, we may replace \tilde{g} by $\tilde{g} - g$ and repeat the whole argument. Since the order of \tilde{K} is finite, we can thus construct hypersurfaces U and U' with the claimed properties after finitely many iterations.

To prove assertion (2), we rewrite the claimed inequality so as to allow a calculation in coordinates. Let $h \in K_1$ be an element of minimal order o . Define for a tuple $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{N}^n$ the number

$$|\gamma|_S = \sum_{i \in S} \gamma_i.$$

Set $s = (s_1, \dots, s_n)$. Since U' is compatible with D' , we know that K'_1 is of the form

$$K'_1 = \left(x_1^{\text{ord } I + |s|_S - c!} \prod_{i \notin T} x_i^{s_i} \right) \cdot I'_1.$$

It is straightforward that $x_1^{-c!} \pi(h) \in K'_1$. This implies that

$$\begin{aligned} \text{ord } I'_1 &= \text{ord } K'_1 - (\text{ord } I + |s|_S - c!) - \sum_{i \notin T} s_i \\ &\leq \text{ord } \pi(h) - \text{ord } I - |s|_S - \sum_{i \notin T} s_i. \end{aligned}$$

Hence, it remains to verify that the inequality

$$\text{ord } \pi(h) \leq \text{ord}_{Q_T} \text{in}(h) + \text{ord } I + |s|_S$$

holds. Let h have the following power series expansion with respect to y :

$$h = \sum_{\alpha \in \mathbb{N}^n} c_\alpha y^\alpha.$$

There exists a multi-index $\beta \in \mathbb{N}^n$ such that $c_\beta \neq 0$, $|\beta| = o$, and

$$\sum_{i \geq 2} \beta_i = \text{ord}_{Q_T} \text{in}(h).$$

Since $\text{coeff}_U(J) = \left(\prod_{i \notin T} x_i^{s_i} \right) \cdot I_1$, we know that $\beta_i \geq s_i$ for all indices $i \notin T$. Consequently,

$$|\beta|_S = o - \sum_{i \notin S} \beta_i \leq o - \sum_{i \notin S} s_i = \text{ord } I + |s|_S.$$

Further,

$$\pi(h) = \sum_{\alpha \in \mathbb{N}^n} c_\alpha x_1^{|\alpha|_S} \prod_{i \geq 2} x_i^{\alpha_i}.$$

From this we conclude that

$$\text{ord } \pi(h) \leq |\beta|_S + \sum_{i \geq 2} \beta_j \leq \text{ord}_{Q_T} \text{in}(h) + \text{ord } I + |s|_S.$$

This proves the claimed inequality. \square

5. PROOF OF THE THEOREM

Using the bound for the order of I'_1 established in the proposition, we can now prove the theorem without considering the coefficient ideals of the weak transform J' of J . Instead, we will directly work in R with the coefficient ideals K and K_1 of J .

Proof of the theorem: Let q , u and K_1 be defined as in the proposition. Set $w = \frac{a}{c!}$. Let p^e be the biggest p -th power dividing c and set $m = \frac{c}{p^e}$. Recall that $\text{ord } q \geq w > 1$.

We begin with assertions (1), (2) and (3). Let $f \in J$ be an element of minimal weighted order cw . Let the weighted initial form of f have the expansion $\text{in}_w(f) = \sum_{i \geq 0} F_i z^i$ with $F_i \in \mathbb{K}[x]$. Hence, either $F_i = 0$ or F_i is a homogeneous polynomial of degree $\deg F_i = (c - i)w$ for all indices i . In particular, F_i is 0 for all indices $i > c$.

Let $i < c$ be an index with $F_i \neq 0$. Due to the factorization $K = M \cdot I$, the element F_i has a factorization

$$F_i = \prod_{j=1}^n x_j^{m_j} \cdot G_i$$

with $m_j \geq \frac{c-i}{c!} s_j$ for some polynomial G_i . Consequently,

$$\begin{aligned} \text{ord}_{Q_T} F_i &= \text{ord}_{Q_T} \underbrace{\prod_{j \in T} x_j^{m_j}}_{=0} + \text{ord}_{Q_T} \prod_{j \notin T} x_j^{m_j} \cdot G_i \\ &\leq \deg \prod_{j \notin T} x_j^{m_j} \cdot G_i \\ &\leq \frac{c-i}{c!} \left(\text{ord } I + \sum_{j \notin T} s_j \right). \end{aligned}$$

Set $H = \text{in}(q)$. Consider the expansion $\text{in}_w(f) = \sum_{i \geq 0} \tilde{F}_i u^i$ with $\tilde{F}_i \in \mathbb{K}[x]$. If $\text{ord } q = w$, then

$$\tilde{F}_i = \sum_{i \leq k \leq c} \binom{k}{i} F_k H^{k-i}.$$

On the other hand, if $\text{ord } q > w$, then $\tilde{F}_i = F_i$. Notice that if $\tilde{F}_i \neq 0$ holds for an index $i < c$, then $\tilde{F}_i \frac{c!}{c-i!}$ is the initial form of an element of minimal order of K_1 . Hence, we know by the proposition that

$$(*) \quad \text{ord } \tilde{F}_i > \frac{c-i}{c!} \left(\text{ord } I + \sum_{j \notin T} s_j \right).$$

Consequently, for all indices $i < c$, either $F_i = 0$ or $\tilde{F}_i \neq F_i$ holds.

This implies that $\text{ord } q = w$. Consequently, w is an integer $w \geq 2$ and we have proved (2).

We show that $F_c \neq 0$. Assume the contrary, and let $i < c$ be maximal with $F_i \neq 0$. Then $\tilde{F}_i = F_i$ would hold by the formula for \tilde{F}_i , contradiction. This shows $F_c \neq 0$. Thus, f is z -regular of order c . After multiplication with a constant, we may assume that $F_c = 1$.

Assume that there exists an index $i < c$ that is not divisible by p^e such that $F_i \neq 0$. Let i be maximal with this property. Then it follows that

$$\tilde{F}_i = F_i + \sum_{\substack{i < k < c \\ p^e | k}} \binom{k}{i} F_k H^{k-i} + \sum_{\substack{i < k < c \\ p^e \nmid k}} \binom{k}{i} \underbrace{F_k}_{=0} H^{k-i} = F_i,$$

contradiction. Therefore F_i is 0 for all indices i that are not divisible by p^e . Set

$$F = \binom{c}{p^e}^{-1} F_{c-p^e}$$

and recall that $\binom{c}{p^e} \not\equiv 0 \pmod{p}$ by Lucas' theorem. Since $\tilde{F}_{c-p^e} = \binom{c}{p^e}(F + H^{p^e})$, we get from (*) the inequality

$$(**) \quad \text{ord}_{Q_T}(F + H^{p^e}) > \frac{p^e}{c!} \left(\text{ord } I + \sum_{j \notin T} s_j \right).$$

Next, we establish for all indices $i < m$ the equality

$$F_{ip^e} = \binom{m}{i} F^{m-i}.$$

Notice that the equality holds by definition for $i = m-1$. Let $i < m-1$ be maximal with $F_{ip^e} \neq \binom{m}{i} F^{m-i}$. Then

$$\begin{aligned} \tilde{F}_{ip^e} &= F_{ip^e} + \sum_{i < k \leq m} \underbrace{\binom{kp^e}{ip^e}}_{\equiv \binom{k}{i}} F_{kp^e} H^{p^e(k-i)} \\ &= F_{ip^e} + \sum_{i < k \leq m} \underbrace{\binom{k}{i} \binom{m}{k}}_{= \binom{m}{i} \binom{m-i}{k-i}} F^{m-k} H^{p^e(k-i)} \\ &= F_{ip^e} - \binom{m}{i} F^{m-i} + \binom{m}{i} (F + G^{p^e})^{m-i}. \end{aligned}$$

Together with the inequalities (*) and (**), this implies that

$$\text{ord}_{Q_T} \left(F_{ip^e} - \binom{m}{i} F^{m-i} \right) > (m-i) \frac{c!}{p^e} \left(\text{ord } I + \sum_{j \notin T} s_j \right).$$

But the factorization $K = M \cdot I$ implies that

$$\text{ord}_{x_j} \left(F_{ip^e} - \binom{m}{i} F^{m-i} \right) > (m-i) \frac{c!}{p^e} s_j$$

holds for all indices $j \in T$. It follows that

$$\begin{aligned} \text{ord}_{Q_T} \left(F_{ip^e} - \binom{m}{i} F^{m-i} \right) &\leq \text{ord} \left(F_{ip^e} - \binom{m}{i} F^{m-i} \right) - (m-i) \frac{c!}{p^e} \sum_{j \in T} s_j \\ &\leq (m-i) \frac{c!}{p^e} \left(\text{ord } I + \sum_{j \notin T} s_j \right), \end{aligned}$$

which contradicts the above inequality. Therefore

$$\text{in}_w(f) = (z^{p^e} + F)^m$$

holds.

Let now $h \in J$ be another element of weighted order cw . Let h have the expansion $h = \sum_{i \geq 0} h_i z^i$. Using the same argument as before, we know that h is z -regular of order c . Hence, h_c is a unit. Consider the element $h - h_c(0) \cdot f \in J$. Since this

element is not z -regular of order c , we know that its weighted order is strictly larger than cw . This implies that

$$\text{in}_w(h) = h_c(0) \cdot \text{in}_w(f) = h_c(0) \cdot (z^{p^e} + F)^m.$$

Assume that F is a p^e -th power. Set $z_1 = z + F^{1/p^e}$ and let V_1 be the regular hypersurface $V_1 = V(z_1)$ in $\text{Spec}(R)$. Then it is straightforward to verify that $\text{ord}_{\text{coeff}_{V_1}(J)} > o$. Thus, the hypersurface V would not have had weak maximal contact with J , contradiction. So assertion (3) is shown. But as F is not a p^e -th power we must have $e \geq 1$, thus proving also (1).

We can now easily prove assertion (4). Notice that

$$\deg G \leq \frac{p^e}{c!} \left(\text{ord } I + \sum_{j \notin T} s_j \right).$$

Using inequality (**), this implies that

$$\text{ord}_{Q_T}^{\text{mod } p^e} F \geq \text{ord}_{Q_T}(F + H^{p^e}) > \deg G.$$

This proves (4).

To prove assertion (5), rewrite (4) as

$$\text{ord}_{(x_2, \dots, x_n)} F((x_1, \dots, x_n) + \mathfrak{t}x_1) + H((x_1, \dots, x_n) + \mathfrak{t}x_1)^{p^e} > v.$$

Setting $x_1 = 1$, this is equivalent to

$$\text{ord } F((1, x_2, \dots, x_n) + \mathfrak{t}) + H((1, x_2, \dots, x_n) + \mathfrak{t})^{p^e} > v.$$

Hence,

$$F((1, x_2, \dots, x_n) + \mathfrak{t}) + H((1, x_2, \dots, x_n) + \mathfrak{t})^{p^e} \in (x_2, \dots, x_n)^{v+1}.$$

Set $N = H((1, x_2, \dots, x_n) + \mathfrak{t})$. Since $F = x^r \cdot G$, we get that

$$G((1, x_2, \dots, x_n) + \mathfrak{t}) - \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i} \cdot N^{p^e}(x_2, \dots, x_n) \in (x_2, \dots, x_n)^{v+1}.$$

But since $\deg G((1, x_2, \dots, x_n) + \mathfrak{t}) \leq v$, this implies that

$$G((1, x_2, \dots, x_n) + \mathfrak{t}) = \lfloor \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i} \cdot N^{p^e}(x_2, \dots, x_n) \rfloor_v$$

as claimed.

We proceed with assertion (6). Assume, by the contrary, that the equality

$$\sum_{i \in T} \bar{r}_i + \bar{v} = b \cdot p^{\ell+1}$$

holds. Computation gives

$$\prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i} = \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-\bar{r}_i} \cdot L^{p^{\ell+1}}(x_2, \dots, x_n)$$

for some element $L \in \mathbb{K}[[x_2, \dots, x_n]]$. Notice that

$$\begin{aligned} \deg \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-\bar{r}_i} &\leq \sum_{i \in T} -\bar{r}_i \\ &= b \cdot p^{\ell+1} - \sum_{i \in T} \bar{r}_i = \bar{v}. \end{aligned}$$

By (5) this implies that

$$G((1, x_2, \dots, x_n) + \mathfrak{t}) = \prod_{i \in T \setminus \{1\}} (x_i + t_i)^{-r_i} \cdot \tilde{N}^{p^{\ell+1}}(x_2, \dots, x_n)$$

for some polynomial \tilde{N} . Consequently,

$$\prod_{i \in T \setminus \{1\}} (x_i + t_i)^{r_i} \cdot G((1, x_2, \dots, x_n) + \mathfrak{t})$$

is a $p^{\ell+1}$ -th power. Since the degree of F is divisible by $p^{\ell+1}$, also $F = x^r \cdot G$ is a $p^{\ell+1}$ -th power, which contradicts the minimality of p^ℓ and proves (6).

It remains to prove assertions (7), (8) and (9). Let k be an integer in the range $0 \leq k < e$. Let $i \in \{1, \dots, n\}$ be an index and assume that

$$\partial_{x_i^{p^k}}(F) \neq 0.$$

Notice that $\partial_{x_i^{p^k}}(H^{p^e}) = 0$. Consequently,

$$\text{ord}_{Q_T} \tilde{F}_{c-p^e} = \text{ord}_{Q_T}(F + H^{p^e}) \leq \text{ord}_{Q_T} \partial_{x_i^{p^k}}(F) + p^k.$$

If $i \in T$, then

$$\partial_{x_i^{p^k}}(F) = x_i^{r_i - p^k} \prod_{\substack{j \in T \\ j \neq i}} x_j^{r_j} H_{i,k}$$

holds for a homogeneous polynomial $H_{i,k} \in \mathbb{K}[x]$ with $\deg H_{i,k} = \deg G$. On the other hand, if $i \notin T$, then

$$\partial_{x_i^{p^k}}(F) = \prod_{j \in T} x_j^{r_j} H_{i,k}$$

for a homogeneous polynomial $H_{i,k} \in \mathbb{K}[x]$ with $\deg H_{i,k} = \deg G - p^k$. Together,

$$\begin{aligned} \text{ord}_{Q_T} \tilde{F}_{c-p^e} &\leq \text{ord}_{Q_T} H_{i,k} + p^k \\ &\leq \deg H_{i,k} + p^k = \deg G + \varepsilon_{i,k} \end{aligned}$$

where

$$\varepsilon_{i,k} = \begin{cases} p^k & \text{if } i \in T, \\ 0 & \text{if } i \notin T. \end{cases}$$

This proves the following inequality:

$$\begin{aligned} \text{ord } I'_1 &\leq \frac{c!}{p^e} \text{ord}_{Q_T} \tilde{F}_{c-p^e} - \sum_{i \notin T} s_i \\ &\leq \frac{c!}{p^e} (\deg G + \varepsilon_{i,k}) - \sum_{i \notin T} s_i \\ &\leq \text{ord } I + \frac{c!}{p^e} \varepsilon_{i,k}. \end{aligned}$$

If $i \notin T$, this implies that $\text{ord } I'_1 \leq \text{ord } I$. Therefore,

$$\partial_{x_i^{p^k}}(F) = 0$$

holds for all indices $i \notin T$ and all $k \geq 0$ with $p^k < p^e$. Thus, the variables x_i with $i \notin T$ only appear as p^e -th powers in F . This proves (7).

Let now ℓ be maximal such that F is a p^ℓ -th power. It is clear that this implies the existence of an index $i \in T$ such that $\partial_{x_i^{p^\ell}}(F) \neq 0$. The inequality above shows

$$\begin{aligned} \text{ord } I'_1 &\leq \text{ord } I + \frac{c!}{p^e} p^\ell \\ &\leq \text{ord } I + \frac{c!}{p}, \end{aligned}$$

which proves (9).

Set $H_i = H_{i,\ell}$ for $i \in T$. Since $\text{ord } I'_1 > \text{ord } I$ and F is a p^ℓ -th power, above inequality also implies for all indices $i \in T$ that

$$\text{ord}_{Q_T} H_i = \text{deg } H_i.$$

This gives (8) and ends the proof of the theorem. □

REFERENCES

- [Abh56] S. Abhyankar, *Local uniformization on algebraic surfaces over ground fields of characteristic $p \neq 0$* , Ann. of Math. (2) **63** (1956), 491–526.
- [Abh66] ———, *Resolution of singularities of embedded algebraic surfaces*, Pure and Applied Mathematics, Vol. 24, Academic Press, New York-London, 1966.
- [BM97] E. Bierstone and P. Milman, *Canonical desingularization in characteristic zero by blowing up the maximum strata of a local invariant*, Invent. Math. **128** (1997), no. 2, 207–302.
- [Cos87] V. Cossart, *Polyèdre caractéristique d'une singularité*, Thèse d'état, Orsay, 1987.
- [Cos91] ———, *Contact maximal en caractéristique positive et petite multiplicité*, Duke Math. J. **63** (1991), no. 1, 57–64.
- [CP08] V. Cossart and O. Piltant, *Resolution of singularities of threefolds in positive characteristic. I. Reduction to local uniformization on Artin-Schreier and purely inseparable coverings*, J. Algebra **320** (2008), no. 3, 1051–1082.
- [CP09] ———, *Resolution of singularities of threefolds in positive characteristic. II*, J. Algebra **321** (2009), no. 7, 1836–1976.
- [Cut04] S. D. Cutkosky, *Resolution of singularities*, American Mathematical Society, Providence, R.I., 2004.
- [Cut09] ———, *Resolution of singularities for 3-folds in positive characteristic*, Amer. J. Math. **131** (2009), no. 1, 59–127.
- [Cut11] ———, *A skeleton key to Abhyankar's proof of embedded resolution of characteristic p surfaces*, Asian J. Math. **15** (2011), no. 3, 369–416.
- [EH02] S. Encinas and H. Hauser, *Strong resolution of singularities in characteristic zero*, Comment. Math. Helv. **77** (2002), no. 4, 821–845.
- [Gir75] J. Giraud, *Contact maximal en caractéristique positive*, Ann. Sci. École Norm. Sup. (4) **8** (1975), no. 2, 201–234.
- [Hau03] H. Hauser, *The Hironaka theorem on resolution of singularities (or: A proof we always wanted to understand)*, Bull. Amer. Math. Soc. (N.S.) **40** (2003), no. 3, 323–403 (electronic).
- [Hau04] ———, *Three power series techniques*, Proc. London Math. Soc. (3) **89** (2004), no. 1, 1–24.
- [Hau10] ———, *On the problem of resolution of singularities in positive characteristic (or: a proof we are still waiting for)*, Bull. Amer. Math. Soc. (N.S.) **47** (2010), no. 1, 1–30.
- [Hau14] ———, *Blowups and resolution*, The resolution of singular algebraic varieties, Amer. Math. Soc., Providence, RI, 2014, pp. 1–80.
- [Hir64] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II*, Ann. of Math. (2) **79** (1964), 109–203; *ibid.* (2) **79** (1964), 205–326.
- [Hir84] ———, *Desingularization of excellent surfaces, Bowdoin 1967*, Resolution of surface singularities, V. Cossart, J. Giraud, and U. Orbanz, Lecture Notes in Mathematics, vol. 1101, Springer-Verlag, 1984.

- [Hir12] ———, *Resolution of singularities*, Manuscript distributed at the CMI Summer School 2012, 138 pp.
- [HP16a] H. Hauser and S. Perlega, *A new proof for the embedded resolution of surface singularities*, Manuscript, 2016.
- [HP16b] ———, *Indefinite increase of the residual order for the resolution of singularities in positive characteristic*, Manuscript, 2016.
- [HW14] H. Hauser and D. Wagner, *Alternative invariants for the embedded resolution of purely inseparable surface singularities*, Enseign. Math. **60** (2014), no. 1-2, 177–224.
- [Kaw07] H. Kawanoue, *Toward resolution of singularities over a field of positive characteristic. I. Foundation; the language of the idealistic filtration*, Publ. Res. Inst. Math. Sci. **43** (2007), no. 3, 819–909.
- [KM10] H. Kawanoue and K. Matsuki, *Toward resolution of singularities over a field of positive characteristic (the idealistic filtration program) Part II. Basic invariants associated to the idealistic filtration and their properties*, Publ. Res. Inst. Math. Sci. **46** (2010), no. 2, 359–422.
- [Kol07] J. Kollár, *Lectures on resolution of singularities*, Princeton University Press, Princeton, 2007.
- [Moh87] T.-T. Moh, *On a stability theorem for local uniformization in characteristic p* , Publ. Res. Inst. Math. Sci. **23** (1987), no. 6, 965–973.
- [Moh96] T. T. Moh, *On a Newton polygon approach to the uniformization of singularities of characteristic p* , Algebraic geometry and singularities (La Rábida, 1991), Progr. Math., vol. 134, Birkhäuser, Basel, 1996, pp. 49–93.
- [Vil89] O. Villamayor, *Constructiveness of Hironaka’s resolution*, Ann. Sci. École Norm. Sup. (4) **22** (1989), no. 1, 1–32.
- [Vil92] ———, *Patching local uniformizations*, Ann. Sci. École Norm. Sup. (4) **25** (1992), no. 6, 629–677.
- [Vil07] ———, *Hypersurface singularities in positive characteristic*, Adv. Math. **213** (2007), no. 2, 687–733.
- [Wlo05] J. Włodarczyk, *Simple Hironaka resolution in characteristic zero*, J. Amer. Math. Soc. **18** (2005), no. 4, 779–822 (electronic).

Faculty of Mathematics
University of Vienna, Austria
herwig.hauser@univie.ac.at
stefan.perlega@univie.ac.at