

# Towards a classification of static electrovacuum spacetimes containing an asymptotically flat spacelike hypersurface with compact interior

Piotr T Chruściel<sup>†</sup>

Département de Mathématiques, Faculté des Sciences, Parc de Grandmont, F37200 Tours, France

Received 6 October 1998

**Abstract.** We show that static electrovacuum black hole spacetimes containing an asymptotically flat spacelike hypersurface with compact interior and with both degenerate and non-degenerate components of the event horizon do not exist, under the supplementary hypothesis that all degenerate components of the event horizon have charges of the same sign. This extends previous uniqueness theorems of Simon and Masood-ul-Alam (where only non-degenerate horizons were allowed) and Heusler (where only degenerate horizons were allowed).

PACS number: 0470B

## 1. Introduction

A classical question in general relativity, first raised and partially answered by Israel [22], is that of classification of non-singular black hole solutions of the Einstein–Maxwell equations. The most complete results existing in the literature so far are due to Simon [28], Masood-ul-Alam [24] and Heusler [19, 21] who show, roughly speaking, the following:

1. Suppose that all the horizons are non-degenerate. Then the black hole is a Reissner–Nordström black hole [19, 24, 28].
2. Suppose instead that all the horizons are degenerate, and that

$$\forall i, j \quad Q_i Q_j \geq 0, \quad (1.1)$$

where  $Q_i$  is the charge of the  $i$ th connected component of the black hole. Then the black hole is a standard Majumdar–Papapetrou black hole [21] (cf also [9, 11]).

Condition (1.1) is obviously satisfied by a connected black hole, so that the above results settle the classification question in the connected case (recall that a standard connected Majumdar–Papapetrou black hole is an extreme Reissner–Nordström one). The general case, however, still remains open. In this paper we ‘merge’ those two results and show the following.

**Theorem 1.1.** *Let  $(M, g, F)$  be a static solution of the Einstein–Maxwell equations with defining Killing vector  $X$ . Suppose that  $M$  contains a connected and simply connected spacelike hypersurface  $\Sigma$  the closure  $\bar{\Sigma}$  of which is the union of an asymptotically flat end and of a compact interior, such that:*

1. *We have  $g_{\mu\nu} X^\mu X^\nu < 0$  on  $\bar{\Sigma}$ .*

<sup>†</sup> Alexander von Humboldt fellow. Supported in part by a grant from the Polish Committee for Scientific Research (KBN) no 2 PO3B 073 15 and by the Humboldt Foundation. E-mail address: [Chrusciel@Univ-Tours.Fr](mailto:Chrusciel@Univ-Tours.Fr)

<sup>‡</sup> We use the signature  $(-, +, +, +)$ .

2. The topological boundary  $\partial\Sigma \equiv \bar{\Sigma} \setminus \Sigma$  of  $\Sigma$  is a non-empty topological manifold, with  $g_{\mu\nu}X^\mu X^\nu = 0$  on  $\partial\Sigma$ .

Then:

1. If  $\partial\Sigma$  is connected, then  $\Sigma$  is diffeomorphic to  $\mathbb{R}^3$  minus a ball. Moreover, there exists a neighbourhood of  $\Sigma$  in  $M$  which is isometrically diffeomorphic to an open subset of the (extreme or non-extreme) Reissner–Nordström spacetime.
2. If  $\partial\Sigma$  is not connected and if condition (1.1) holds for charges  $Q_i$  associated with those components of  $\partial\Sigma$  that intersect the degenerate horizons, then  $\Sigma$  is diffeomorphic to  $\mathbb{R}^3$  minus a finite union of disjoint balls. Moreover, the spacetime contains only degenerate horizons, and there exists a neighbourhood of  $\Sigma$  in  $M$  which is isometrically diffeomorphic to an open subset of the standard Majumdar–Papapetrou spacetime.

Actually a somewhat more general result is proved in theorem 3.6 below. We emphasize that no sign conditions are made concerning the charges of non-degenerate horizons. We also note that simple connectedness of  $\Sigma$  will hold when appropriate further global hypotheses on  $M$  are made, cf theorem 1.3 below. Thus, to obtain a satisfactory classification of the spacetimes under consideration it remains to remove the condition on the sign of the charges, or to construct (and classify) appropriately regular black holes which do not satisfy this condition. We find that last possibility rather unlikely.

The definitions and conventions used here coincide with those of the accompanying paper [10]. Those definitions which cannot be found there are presented in section 2 below.

We refer the reader to a discussion of a similar theorem for vacuum spacetimes in section 1 of [10] for comments concerning the improvements of this result compared to those available in the literature even in cases where a mixture of degenerate and non-degenerate horizons is forbidden. It might be of some interest to mention that our conclusion will still hold for a somewhat larger class of manifolds  $\Sigma$ . A possible generalization is that with  $\Sigma$  being, for example, the union of (a) a finite number of asymptotically flat ends with (b) a neighbourhood of the boundary  $\partial\Sigma$  which has compact closure in  $M$  and (c) a non-compact region on which we have  $0 < \epsilon \leq 1 + \phi \pm \sqrt{-g_{\mu\nu}X^\mu X^\nu}$ , provided that  $\Sigma$  with the induced metric is a *complete* Riemannian manifold;  $\phi$  here is the electric potential as defined in equation (3.3) after the relevant duality rotations have been performed, cf lemma 3.2. The proof carries through without any modifications to this case.

Our strategy is a modification of that of Ruback† [27] along the lines of [10]: we consider the *orbit space* metric  $h$  on  $\Sigma$ , as defined in [10]. The key tools here are the results of [10] concerning the geometry of  $(\Sigma, h)$  near both the degenerate components of  $\partial\Sigma$  and the non-degenerate ones. Next, following [27], we consider a manifold which consists of two copies of  $(\Sigma, h)$  glued along all *non-degenerate* components of  $\partial\Sigma$ , equipped with an appropriate conformally deformed metric. As in [10] we use a new version of the positive energy theorem proved in [3] (theorem 3.3 below) to show that the metric on  $\Sigma$  is conformally flat. One can then use classical calculations to finish the proof. We note that it is usual in the last step of the proof to invoke analyticity to conclude. Because analytic extensions of manifolds are not unique this is not sufficient without a more thorough justification. We finish the proof by a simple open–closed argument which avoids this problem.

† We note that while the relevant claims in [27] can eventually be justified, this reference contains several essential gaps. The work here can be considered as an extension of that of Ruback to include degenerate black holes, together with a justification of the relevant unsubstantiated claims made in [27]. We further note that we have not been able to adapt the technique of Simon [28] and Masood-ul-Alam [24] to include degenerate black holes without having to introduce some supplementary restrictions.

Under the hypotheses of theorem 1.1 there is no chance of getting more information about the size of the set on which the metric is that of a Reissner–Nordström or a standard Majumdar–Papapetrou spacetime (consider any hypersurface  $\Sigma$  in the Reissner–Nordström spacetime, and set  $M$  to be any neighbourhood of  $\Sigma$  which does not coincide with the Reissner–Nordström spacetime; alternatively, identify  $t$  with  $t+1$  in the Reissner–Nordström spacetime). In complete analogy with the vacuum case in [10] we have the following.

**Corollary 1.2.** *Under the hypotheses of theorem 1.1, assume further that*

3. *The orbits of the Killing vector  $X$  through  $\Sigma$  are complete.*

*Then the following properties are equivalent:*

- (i)  $\Sigma_{\text{ext}}$  is achronal<sup>†</sup> in  $M_{\text{ext}}$ .
- (ii)  $M_{\text{ext}}$  is diffeomorphic to  $\mathbb{R} \times \Sigma_{\text{ext}}$  (which is equivalent to an appropriately complete  $\mathcal{J}$  having  $\mathbb{R} \times S^2$  topology).
- (iii) *There are no closed timelike curves through  $\Sigma_{\text{ext}}$  contained in  $M_{\text{ext}}$ .*

*Further, if one (and hence all) of the above conditions holds, then the Killing development<sup>‡</sup>  $\mathcal{K}(\Sigma)$  of  $\Sigma$  defined as*

$$\mathcal{K}(\Sigma) \equiv \cup_{t \in \mathbb{R}} \phi_t(\Sigma), \quad (1.2)$$

*where  $\phi_t$  is the action of the isometry group generated by  $X$ , equipped with the induced metric, is isometrically diffeomorphic to a domain of outer communications of a standard extension of a Reissner–Nordström spacetime or of a standard Majumdar–Papapetrou spacetime.*

The *standard Majumdar–Papapetrou* spacetimes are defined in section 2. We refer the reader to the introduction of [10] for a discussion of the relationship between theorem 1.1 and corollary 1.2 and black holes. In particular, in the introduction of [10] an example was given which shows that more hypotheses than those of corollary 1.2 are needed to show that  $\mathcal{K}(\Sigma)$  coincides with a d.o.c. in  $M$ . For reference we state the following.

**Theorem 1.3.** *Let  $(M, g, F)$  be a solution of the Einstein–Maxwell equations containing a connected spacelike hypersurface  $\Sigma$ , the closure  $\bar{\Sigma}$  of which is the union of a finite number of asymptotically flat ends and of a compact interior. Let  $X$  be a Killing vector field on  $M$  which is timelike future directed in all the asymptotically flat ends, and which satisfies the staticity condition (2.1). Further, let  $\mathcal{D}_{oc} \equiv \mathcal{D}_{oc}(M_{\text{ext}})$  be a domain of outer communications in  $(M, g)$  associated with one of the asymptotically flat ends of  $\Sigma$ . Suppose that:*

1. *We have  $\Sigma \subset \mathcal{D}_{oc}$ .*
2. *The topological boundary  $\partial \Sigma \equiv \bar{\Sigma} \setminus \Sigma$  of  $\Sigma$  is a non-empty topological manifold and satisfies  $\partial \Sigma = \bar{\Sigma} \cap \partial \mathcal{D}_{oc}$ .*
3.  *$X$  has complete orbits in  $\mathcal{D}_{oc}$ .*

*In addition to the above, suppose that condition (1.1) holds for charges  $Q_i$  associated to those components of  $\partial \Sigma$  that intersect the degenerate horizons and that one of the following conditions holds:*

- 4(a) *Either  $(\mathcal{D}_{oc}, g|_{\mathcal{D}_{oc}})$  is globally hyperbolic, or*
- 4(b)  *$(M, g)$  is globally hyperbolic.*

<sup>†</sup> By that we mean that there are no timelike curves from  $\Sigma_{\text{ext}}$  to itself which are entirely contained in  $M_{\text{ext}}$ .

<sup>‡</sup> The notion of Killing development used here differs slightly from the definition given in [5], as here we allow a topology of  $\mathcal{K}(\Sigma)$  which is not  $\mathbb{R} \times \Sigma$ .

Then the conclusions of theorem 1.1 and corollary 1.2 hold. Moreover,  $\mathcal{D}_{oc}$  is isometrically diffeomorphic to a domain of outer communications of a standard extension of a Reissner–Nordström spacetime or of a standard Majumdar–Papapetrou spacetime.

We note that it is not assumed above that  $X$  is timelike throughout  $\Sigma$ .

The proofs of both corollary 1.2 and theorem 1.3 are essentially identical to the corresponding ones in [10]; some comments about the proof of corollary 1.2 can be found at the end of section 3; the proof of theorem 1.3 will be omitted. We note that the property that  $\Sigma$  is simply connected and has only one asymptotically flat end required in theorem 1.1 follows from [12]. We further note that the obvious electrovacuum generalization of the remaining cases of theorem 1.3 of [10] holds under the supplementary hypothesis that  $\Sigma$  is simply connected and has only one asymptotically flat end.

This paper is organized as follows. Section 2 contains definitions and some preliminary remarks. In section 3 we prove theorem 1.1, as a consequence of the somewhat more general theorem 3.6, which is also proved there.

## 2. Preliminaries

Our conventions and definitions are as in section 2 of [10]. Further, a triple  $(M, g, F)$  will be said to be static if there exists on  $M$  a Killing vector field  $X$  such that the Maxwell 2-form field  $F$  satisfies

$$\mathcal{L}_X F = 0,$$

with  $X$  satisfying moreover the staticity condition

$$X_{[\alpha} \nabla_{\beta} X_{\gamma]} = 0. \quad (2.1)$$

Here and throughout  $\mathcal{L}_X$  denotes the Lie derivative with respect to  $X$ .

Next, a data set  $(\Sigma_{\text{ext}}, g, K)$  with Maxwell field  $F$  will be called an *asymptotically flat end* if  $\Sigma_{\text{ext}}$  is diffeomorphic to  $\mathbb{R}^3$  minus a ball and if the fields  $(g_{ij}, K_{ij})$  satisfy the fall-off conditions

$$|g_{ij} - \delta_{ij}| + r|\partial_\ell g_{ij}| + \cdots + r^k |\partial_{\ell_1 \dots \ell_k} g_{ij}| + r|K_{ij}| + \cdots + r^k |\partial_{\ell_1 \dots \ell_{k-1}} K_{ij}| \leq C_{k,\alpha} r^{-\alpha}, \quad (2.2)$$

for some constants  $C_{k,\alpha}$ ,  $\alpha > 0$ ,  $k \geq 1$ . We shall further require that in the local coordinates as above on  $\Sigma_{\text{ext}}$  the Maxwell field satisfies the fall-off conditions

$$|F_{\mu\nu}| + r|\partial_\ell F_{\mu\nu}| + \cdots + r^k |\partial_{\ell_1 \dots \ell_k} F_{\mu\nu}| \leq \hat{C}_{k,\alpha} r^{-\alpha-1}, \quad (2.3)$$

for some constants  $\hat{C}_{k,\alpha}$ ,  $\alpha > 0$ ,  $k \geq 0$ . We shall always implicitly assume  $\alpha > \frac{1}{2}$  when the ADM mass will be invoked, as this condition makes it well defined in vacuum. It follows in any case from section 1.3 of [8] that in stationary electrovacuum spacetimes there is no loss of generality in assuming  $\alpha = 1$ ,  $k$  arbitrary. A hypersurface will be said to be *asymptotically flat* if it contains an asymptotically flat end  $\Sigma_{\text{ext}}$ .

To avoid ambiguities, we define the *Reissner–Nordström* spacetime  $(M^{\text{RN}}, g^{\text{RN}})$  to be the manifold  $\{t \in \mathbb{R}, r \in (m + \sqrt{m^2 - Q^2 - P^2}, \infty), q \in S^2\}$ , with

$$m^2 - Q^2 - P^2 \geq 0, \quad (2.4)$$

and with the metric

$$g^{\text{RN}} = -\left(1 - \frac{2m}{r} + \frac{Q^2 + P^2}{r^2}\right) dt^2 + \left(1 - \frac{2m}{r} + \frac{Q^2 + P^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2, \quad (2.5)$$

where  $d\Omega^2$  is the standard round metric on a unit two-dimensional sphere  $S^2$ . It is somewhat awkward to build in the inequality (2.4) in our definition of a Reissner–Nordström spacetime,

but it saves us the need of repeating that (2.4) holds each time we mention a Reissner–Nordström spacetime. The Maxwell field is

$$F \equiv F_{\mu\nu} dx^\mu \wedge dx^\nu = \frac{Q}{r^2} dt \wedge dr - P \sin(\theta) d\theta \wedge d\phi, \tag{2.6}$$

so that  $Q$  is the total electric charge and  $P$  is the total magnetic charge of  $\Sigma_{\text{ext}}$ . We will refer to those coordinates as the standard coordinates on the Reissner–Nordström spacetime. We shall call the *standard extension of the Reissner–Nordström spacetime* the extension of  $(M^{\text{RN}}, g^{\text{RN}})$  described, for example, by the Carter–Penrose diagram on p 158 of [17] for  $m^2 > Q^2 - P^2$  and on p 160 of [17] for  $m^2 = Q^2 - P^2$ .

Recall that the Majumdar–Papapetrou (MP) metrics are, locally, of the form [23, 26]

$$g = -u^{-2} dt^2 + u^2(dx^2 + dy^2 + dz^2), \tag{2.7}$$

$$A = u^{-1} dt, \tag{2.8}$$

where  $A$  is the Maxwell potential,  $F = dA$ , with some nowhere vanishing, say positive, function  $u$ . A spacetime will be called a standard MP spacetime if the coordinates  $x^\mu$  of (2.7) and (2.8) are global with range  $\mathbb{R} \times (\mathbb{R}^3 \setminus \{\vec{a}_i\})$  for a finite set of points  $\vec{a}_i \in \mathbb{R}^3, i = 1, \dots, I$ , and if the function  $u$  has the form

$$u = 1 + \sum_{i=1}^I \frac{m_i}{|\vec{x} - \vec{a}_i|}, \tag{2.9}$$

for some positive constants  $m_i$ . It has been shown by Hartle and Hawking [16] that every standard MP spacetime can be analytically extended to an electrovacuum spacetime with a non-empty black hole region, and with a domain of outer communication which is non-singular in the sense of the theorems proved here. Those extensions will be called the *standard extensions of the standard Majumdar–Papapetrou spacetimes*.

### 3. Proof of theorem 1.1

Following [10], we equip  $\Sigma$  with the orbit space metric  $h$  defined as

$$h(Y, Z) = g(Y, Z) - \frac{g(X, Y)g(X, Z)}{g(X, X)}, \tag{3.1}$$

where  $X$  is the defining Killing vector, that is, the Killing vector which asymptotes  $\partial/\partial t$  in the asymptotic regions, and satisfies the staticity condition (2.1). Let the electric field  $E$  and the magnetic field  $B$  be defined on  $M$  by the equations (we use the conventions of [20])

$$E(Y) = -F(X, Y), \quad B(Y) = (*F)(X, Y), \tag{3.2}$$

where  $*F$  denotes the spacetime Hodge dual of the Maxwell field 2-form  $F$ . Simple connectedness of  $\Sigma$  and a standard calculation (cf, for example, [20]) shows that there exist functions  $\phi$  and  $\psi$  defined in a neighbourhood of  $\Sigma$  in  $M$  such that we have

$$E = d\phi, \quad B = d\psi, \quad \mathcal{L}_X \phi = \mathcal{L}_X \psi = 0. \tag{3.3}$$

By an abuse of notation we shall often use the symbol  $\phi$  to denote the restriction of  $\phi$  to  $\Sigma$ , similarly with  $\psi$ . The potentials  $\phi$  and  $\psi$  are of course defined up to a constant, and we can normalize them so that on  $\Sigma_{\text{ext}}$  we have

$$\phi = \frac{Q}{r} + O(r^{-2}), \quad \psi = \frac{P}{r} + O(r^{-2}), \tag{3.4}$$

where  $Q$  is the total electric charge and  $P$  is the total magnetic charge in  $\Sigma_{\text{ext}}$ . (We note if there were several asymptotically flat ends it could happen that the potentials could asymptote

constants different from zero on some ends, and the proof given below would break down. This is the only place where the hypothesis that  $\Sigma$  has only one end enters in the argument. In fact, one could allow several ends when the supplementary hypothesis is made that  $\phi$  and  $\psi$  can be normalized to asymptote to zero in all asymptotically flat ends.)

The metric  $h$  on  $\Sigma$  is essentially ‘the metric that would have been induced on  $\Sigma$  if  $\Sigma$  were normal to  $X$ ’, so that we have the following equivalent of lemma 5.1 of [10], the proof of which is a repetition of that in [10].

**Lemma 3.1.** *Suppose that  $(M, g, F)$  is static and assume that the set  $(\hat{h}, \hat{V}, \hat{\phi})$ , where  $\hat{h}$  is the metric induced on the hypersurfaces orthogonal to  $X$ ,  $-\hat{V}^2$  is the square of the Lorentzian norm of  $X$  on those hypersurfaces, and  $\hat{\phi}$  (respectively  $\hat{\psi}$ ) is the restriction of the electric potential  $\phi$  (respectively the magnetic potential  $\psi$ ) defined by equations (3.2) and (3.3) to those hypersurfaces, satisfies some coordinate-independent system of equations. Then the orbit space-metric  $h$  together with the function  $V$  (such that  $-V^2$  is the square of the Lorentzian norm of  $X$  on  $\Sigma$ ) and the electric potential  $\phi|_{\Sigma}$  and the magnetic potential  $\psi|_{\Sigma}$  satisfy the same system of equations.*

It follows that in the Einstein–Maxwell case we have the equations:

$$V \Delta_h \phi = h(d\phi, dV), \quad (3.5)$$

$$V \Delta_h \psi = h(d\psi, dV), \quad (3.6)$$

$$V \Delta_h V = h(d\phi, d\phi) + h(d\psi, d\psi), \quad (3.7)$$

$$V R_{ij} = D_i D_j V + V^{-1} \{ (h(d\phi, d\phi) + h(d\psi, d\psi)) h_{ij} - 2\phi_{,i} \phi_{,j} - 2\psi_{,i} \psi_{,j} \}, \quad (3.8)$$

where  $\Delta_h$  is the Laplace operator of the metric  $h$ ,  $R_{ij}$  is the Ricci tensor of  $h$  and where a comma denotes differentiation. In particular, we have

$$R \equiv h^{ij} R_{ij} = 2V^{-2} \{ h(d\phi, d\phi) + h(d\psi, d\psi) \}. \quad (3.9)$$

Following Heusler [19] we note:

**Lemma 3.2.** *Under the hypotheses of theorem 1.1 the magnetic field  $B$  can be made to vanish by a duality rotation.*

**Proof.** If  $E \equiv 0$  the result is obvious by exchanging  $\phi$  with  $\psi$ . Suppose thus that  $E$  is not identically vanishing, as shown, for example, in [19] we then have

$$B = \mu E,$$

where  $\mu$  is constant on each connected component of the set  $\Omega \equiv \{E \neq 0\}$ . Let  $\Omega_0$  be any connected component of  $\Omega$ , by performing a duality rotation we can obtain  $\psi = 0$  in  $\Omega_0$  [19]. As  $\Omega_0$  is open, equation (3.6) and the unique continuation theorem of Aronszajn [1] show that  $\psi \equiv 0$ , hence  $B \equiv 0$ .  $\square$

Unless explicitly stated otherwise, in the remainder of the paper we shall assume that the duality transformation of lemma 3.2 has been performed, so that

$$\psi \equiv 0.$$

In the proof of theorem 1.1 we shall need the following version of the positive energy theorem, proved in [3]:

**Theorem 3.3.** Let  $(\hat{\Sigma}, \hat{h})$  be a smooth complete Riemannian manifold with an asymptotically flat end  $\hat{\Sigma}_{\text{ext}}$  (in the sense of equation (2.2) with  $k \geq 4$  and  $\alpha > \frac{1}{2}$ ) and with a smooth divergence-free vector field  $\hat{E}$  satisfying

$$\hat{E}_i dx^i = \frac{\hat{Q}}{r^2} dr + o(r^{-2})$$

in  $\hat{\Sigma}_{\text{ext}}$ . Suppose that the Ricci scalar  $\hat{R}$  of  $\hat{h}$  satisfies

$$0 \leq \hat{R} - 2\hat{h}(\hat{E}, \hat{E}) \in L^1(\hat{\Sigma}_{\text{ext}}).$$

Then the ADM mass  $\hat{m}$  of  $\hat{\Sigma}_{\text{ext}}$  satisfies

$$\hat{m} \geq |\hat{Q}|,$$

where  $\hat{Q}$  is the total charge of  $\hat{\Sigma}_{\text{ext}}$ . If the equality is attained and  $\hat{E}$  is not identically vanishing, then the metric  $\hat{h}$  is, locally, the metric induced on the  $t = \text{constant}$  slices of a Majumdar–Papapetrou spacetime (cf equation (2.7)) with

$$\hat{E}_i dx^i = \frac{du}{u}, \quad (3.10)$$

where  $u$  is as in (2.7) and (2.8).

We emphasize that in the result above  $\hat{\Sigma}$  can have an arbitrary number (perhaps infinite) of asymptotic ends, and that *no hypotheses* are made on the asymptotic behaviour of the metric in those ends except that the metric  $\hat{h}$  is complete (and that at least one of the ends is asymptotically flat so that its ADM mass is well defined). More general results, allowing for non-vanishing extrinsic curvature of the initial data hypersurface, non-vanishing of the magnetic field, poor differentiability of the metric, and boundaries, can be found in [3]. The proof of theorem 3.3 uses a Witten-type spinorial argument based on a suggestion of Gibbons and Hull [13]. The equality case is handled by the results of Tod [30]; the plane waves case allowed by Tod is excluded by theorem 3.4 of [5]. We note that in the  $\hat{m} = |\hat{Q}|$  case it is not known whether one can conclude that the metric must be (locally or globally) a standard Majumdar–Papapetrou metric.

To proceed further, we need to analyse the behaviour of  $h$  and  $\phi$  near  $\partial\Sigma$ . Here we shall give an overview of the results needed, and we refer the reader to [10] for detailed proofs of the results discussed in this paragraph. Recall, thus, that by the Vishveshwara–Carter lemma [6, 31]  $\partial\Sigma$  must be a subset of (the closure of) a Killing horizon  $\bar{\mathcal{N}}$ . By that same lemma one knows that in a static spacetime the Killing horizon is a smooth submanifold. Standard results [20] show that  $\phi$  is constant on any connected component of  $\bar{\mathcal{N}}$ , and hence of  $\partial\Sigma$ . A connected component  $S$  of  $\partial\Sigma$  will be called degenerate, respectively non-degenerate, if  $S$  intersects a degenerate, respectively non-degenerate, Killing horizon. By deforming  $\Sigma$  slightly in spacetime if necessary we can ensure that  $\partial\Sigma$  is a smooth submanifold both of  $\bar{\Sigma}$  and of  $M$  near degenerate horizons. Every degenerate component corresponds to a complete end of  $(\Sigma, h)$  (see proposition 3.2 of [10]). As far as non-degenerate horizons are concerned,  $\partial\Sigma$  will not be a smooth submanifold of  $M$  in general when there are points on  $\partial\Sigma$  at which the Killing vector field  $X$  vanishes. However, we can equip  $\bar{\Sigma}$  with a differentiable structure so that  $\partial\Sigma$  is a smooth submanifold of  $\bar{\Sigma}$  (see proposition 3.3 of [10]). Moreover,  $\partial\Sigma$  with this differentiable structure is a totally geodesic boundary of  $(\Sigma, h)$  across which  $h$  can be extended smoothly when doubling  $\Sigma$ . Now  $\phi$  is a smooth function on spacetime, and the proof of proposition 3.3 of [10] shows that  $\phi$  is a smooth function of  $(x^2, y^a)$  (here  $x^2$  denotes the square of  $x$ , and not an index 2 on  $x$ ) in an appropriate coordinate system near a non-degenerate

connected component  $S$  of  $\partial\Sigma$ , with  $S$  given by  $x = 0$  in this coordinate system. This implies, in particular, that

$$|d\phi|_h(x = 0) = 0, \quad (3.11)$$

and that  $\phi$  extends smoothly across  $S$  when a doubling of  $\Sigma$  across  $S$  is performed.

We have the following, which is based on an observation of Ruback [27].

**Proposition 3.4.** *Under the hypotheses of theorem 1.1 we have*

$$0 \leq V + |\phi| \leq 1 \quad (3.12)$$

on  $\bar{\Sigma}$ , with the inequalities being strict on  $\Sigma$  except if the metric is, locally, a Majumdar–Papapetrou metric. Further, the right inequality is strict on non-degenerate horizons.

**Proof.** Set

$$F_{\pm} = V^2 - (1 \pm \phi)^2;$$

as noted by Ruback [27] the functions  $F_{\pm}$  satisfy the equation

$$\Delta_{\gamma} F_{\pm} = 0, \quad (3.13)$$

where  $\Delta_{\gamma}$  is the Laplace operator of the metric  $V^{-2}h_{ij}$ . In the asymptotically flat region of  $\Sigma_{\text{ext}}$  the  $F_{\pm}$ 's approach zero, while at every component of  $\partial\Sigma$  we have  $F_{\pm} \leq 0$ .

Suppose, first, that  $F_- = 0$  on all components of  $\partial\Sigma$ ; the maximum principle implies then

$$F_- \equiv 0$$

on  $\bar{\Sigma}$ . Equation (3.8) and the transformation rule of the Ricci tensor under conformal transformations show that the metric  $(1 - \phi)^{-2}h_{ij}$  is Ricci flat. In dimension three this implies flatness, and the proof of lemma 5.1 of [10] shows that near  $\Sigma$  the spacetime metric can locally be written in the Majumdar–Papapetrou form (2.7). A similar analysis applies if  $F_+$  vanishes throughout  $\partial\Sigma$ .

It remains to consider the case in which both  $F_+$  and  $F_-$  are negative somewhere on  $\partial\Sigma$ . From the maximum principle one obtains

$$F_{\pm} < 0 \quad (3.14)$$

on  $\Sigma$ , so that

$$V^2 < (1 - \phi)^2, \quad V^2 < (1 + \phi)^2 \quad (3.15)$$

on  $\Sigma$ .  $V$  has no zeros on  $\Sigma$  by hypothesis, which together with (3.15) shows that both  $1 - \phi$  and  $1 + \phi$  have no zeros on  $\Sigma$ . As both  $1 - \phi$  and  $1 + \phi$  go to 1 at the infinity of  $\Sigma_{\text{ext}}$  it follows that

$$-1 < \phi < 1 \quad (3.16)$$

on  $\Sigma$ . Equations (3.15) and (3.16) imply  $0 < V < \min(1 + \phi, 1 - \phi) = 1 - |\phi|$  on  $\Sigma$ , as desired.

It remains to consider what happens on non-degenerate components of  $\partial\Sigma$ . Thus, let  $S$  be a connected non-degenerate component of  $\partial\Sigma$ , so that  $d\phi$  vanishes on  $S$  by equation (3.11). It is well known, and in any case easily checked from the formulae in section 3 of [10], that

$$|dV|_h(x = 0) = \kappa, \quad (3.17)$$

where  $\kappa$  is the surface gravity of  $S$ ; the condition that  $S$  is non-degenerate is precisely  $\kappa \neq 0$ . Suppose that  $\phi = 1$  on  $S$ , then  $F_-$  vanishes on  $S$  and equation (3.17) shows that  $F_- = \kappa^2 x^2 + O(x^4)$  will be positive in a neighbourhood of  $S$  (recall that  $\phi - \phi|_S = O(x^2)$ ),

which contradicts (3.14). Similarly  $\phi = -1$  on  $S$  would lead to  $F_+$  being positive in a neighbourhood of  $S$ , again a contradiction.  $\square$

We note the following corollary<sup>†</sup> of proposition 3.4.

**Corollary 3.5.** *Under the hypotheses of theorem 1.1 we have*

$$m \geq |Q|, \tag{3.18}$$

where  $m > 0$  is the ADM mass of  $\Sigma_{\text{ext}}$  and  $Q$  the total charge of  $\Sigma_{\text{ext}}$ . Further, if the inequality is attained the metric is, locally, a Majumdar–Papapetrou metric.

**Proof.** A theorem of Beig [4] (cf also [2, 7]) shows that the Komar mass of a static asymptotically vacuum end  $(\Sigma_{\text{ext}}, g|_{\Sigma_{\text{ext}}})$  coincides with its ADM mass, so that we have

$$V = 1 - \frac{m}{r} + O(r^{-2}). \tag{3.19}$$

The inequality (3.18) follows immediately from proposition 3.4 and the asymptotic expansion (3.4). If  $m = Q$  we have  $F_+ = O(r^{-2})$ , and  $F_+ \equiv 0$  follows from (3.13) and the asymptotic strong maximum principle of [29] (appendix). The conclusion that the metric is locally a Majumdar–Papapetrou metric follows then as in the proof of proposition 3.4. The case  $m = -Q$  follows similarly by considering  $F_-$ . The inequality  $m > 0$  follows either from the asymptotic strong maximum principle of [29] (appendix) or from [18].  $\square$

It follows from proposition 3.4 that  $\phi$  satisfies the inequality  $-1 < \phi < 1$  on  $\Sigma$ , and that the values  $\phi = 1$  or  $-1$  can only be attained at degenerate components of  $\partial\Sigma$ . When only one component of the event horizon is degenerate we can without loss of generality assume, changing  $\phi$  to  $-\phi$  if necessary, that we have

$$-1 < \phi \leq 1 \quad \text{on } \bar{\Sigma}. \tag{3.20}$$

It is tempting to conjecture that one can *always* assume, changing  $\phi$  to  $-\phi$  if necessary, that

$$0 \leq \phi \leq 1 \quad \text{on } \bar{\Sigma}.$$

This is due to the fact that a change of the sign of  $\phi$  will necessarily lead to both positive and negative charges of event horizons, cf lemma 3.7 below—such a configuration is unlikely to be static. Whatever the situation, if equation (3.20) holds we can prove the following.

**Theorem 3.6.** *Let  $(M, g, F)$  be a static solution of the Einstein–Maxwell equations with defining Killing vector  $X$ . Suppose that  $M$  contains a connected and simply connected spacelike hypersurface  $\Sigma$  the closure  $\bar{\Sigma}$  of which is the union of an asymptotically flat end and of a compact interior, such that:*

1. *We have  $g_{\mu\nu}X^\mu X^\nu < 0$  on  $\Sigma$ .*
2. *The topological boundary  $\partial\Sigma \equiv \bar{\Sigma} \setminus \Sigma$  of  $\Sigma$  is a non-empty topological manifold, with  $g_{\mu\nu}X^\mu X^\nu = 0$  on  $\partial\Sigma$ .*

*If equation (3.20) holds, then the conclusions of theorem 1.1 hold.*

<sup>†</sup> The inequality (3.18) has been established under rather more general circumstances in [18] (remark, p 107), using a technique suggested by Gibbons and Hull [13].

**Proof.** The case  $m = |Q| = 0$  cannot occur by [18]. If  $m = |Q| \neq 0$  the metric is, locally, of Majumdar–Papapetrou form by corollary 3.5. In that case we can apply [9] (theorem 7.2; cf also [11]) to the Killing development  $(\hat{M}, \hat{g})$  of  $\Sigma$  as defined in [5] to conclude that  $(\hat{M}, \hat{g})$  is a standard Majumdar–Papapetrou spacetime, and the result follows; cf the argument around equation (3.35) below for a more detailed exposition of the construction of the embedding in the Reissner–Nordström context.

It remains to analyse the case  $m > |Q|$ . In order to do that, consider the manifold  $\Sigma$  equipped with the metric  $h$  defined by equation (3.1). From what has been said  $(\Sigma, h)$  is a complete Riemannian manifold with compact (perhaps empty) boundary and with at least one asymptotically flat end  $\Sigma_{\text{ext}}$ . Let us denote by  $\partial_{nd}\Sigma$  the collection of all those components of the boundary of  $\Sigma$  which correspond to non-degenerate components of the event horizon of the black hole. Following [27], if  $\partial_{nd}\Sigma \neq \emptyset$  we set

$$\begin{aligned}\Sigma_+ &= \Sigma, & h_+ &= \left(\frac{1}{2}(1+V+\phi)\right)^2 h, \\ \Sigma_- &= \Sigma, & h_- &= \left(\frac{1}{2}(1-V+\phi)\right)^2 h,\end{aligned}\tag{3.21}$$

$$\begin{aligned}\hat{\Sigma} &= \Sigma_+ \cup \Sigma_- \cup \partial_{nd}\Sigma, & \hat{h}|_{\Sigma_+} &= h_+, & \hat{h}|_{\Sigma_-} &= h_-, \\ \hat{E}_{\pm} &= \frac{(1+\phi)d\phi - VdV}{V(1+\phi \pm V)}, & \hat{E}|_{\Sigma_+} &= \hat{E}_+, & \hat{E}|_{\Sigma_-} &= \hat{E}_-.\end{aligned}\tag{3.22}$$

The topological and differentiable structure of  $\hat{\Sigma}$  are defined through the gluing of  $\bar{\Sigma}_+ \equiv \Sigma_+ \cup \partial_{nd}\Sigma$  with  $\bar{\Sigma}_- \equiv \Sigma_- \cup \partial_{nd}\Sigma$  by identifying  $\partial_{nd}\Sigma$ , considered as a subset of  $\bar{\Sigma}_+$ , with a second copy of  $\partial_{nd}\Sigma$ , considered as a subset of  $\bar{\Sigma}_-$ , using the identity map. From our remarks at the beginning of this section it follows that the metric  $\hat{h}$  defined on  $\Sigma_+ \cup \Sigma_-$  in (3.21) can be extended by continuity to a smooth metric on  $\hat{\Sigma}$ ; similarly  $\hat{E}$  can be extended by continuity to a smooth vector field on  $\hat{\Sigma}$ .

If  $\partial_{nd}\Sigma = \emptyset$  we set

$$\hat{\Sigma} = \Sigma, \quad \hat{h} = h_+, \quad \hat{E} = E_+.$$

We have the following:

- The conformal factor  $1 - V + \phi \geq 1 - V - |\phi|$  is strictly positive on  $\Sigma \cup \partial_{nd}\Sigma$  by proposition 3.4, and so is  $1 + V + \phi = 2V + 1 - V + \phi \geq 1 - V + \phi$ . Near every connected degenerate component  $S$  of  $\partial\Sigma$  the electric potential  $\phi$  will tend to a value different from  $-1$  by the hypothesis (3.20), while  $V$  will tend to zero, hence the asymptotic end of  $(\Sigma, h)$  corresponding to  $S$  remains complete in the metric  $(\hat{\Sigma}, \hat{h})$ .
- The conformal factor  $(1 + V + \phi)/2$  tends to 1 in the asymptotically flat end  $\Sigma_{\text{ext}}$ , so that  $\Sigma_{\text{ext}}$  is an asymptotically flat end for the metric  $h_+$ , with ADM mass equal to

$$\hat{m} = \frac{1}{2}(m + Q).$$

The electric field  $\hat{E}$  approaches zero as  $1/r^2$  in  $\Sigma_{\text{ext}}$  and has charge equal to

$$\hat{Q} = \frac{1}{2}(m + Q) = \hat{m}.$$

- The conformal factor  $1 - V + \phi$  tends to 0 in the asymptotically flat  $\Sigma_{\text{ext}}$  as  $(m + Q)/r$ , with  $m + Q \neq 0$ , thus as  $r$  tends to infinity in  $\Sigma_{\text{ext}}$  the metric  $h_-$  approaches, to leading significant orders, the metric

$$\frac{1}{4}(m + Q)^2 \left( \frac{1}{r^2} dr^2 + d\Omega^2 \right),$$

where  $d\Omega^2$  is the standard round metric on a 2-sphere. It easily follows that  $(\Sigma_{\text{ext}}, h_-)$  is a *complete* end of  $(\hat{\Sigma}, \hat{h})$  (cf the calculation in the proof of proposition 3.2 in [10]).

As emphasized by Ruback [27] we have

$$\hat{R} = 2\hat{h}(\hat{E}, \hat{E}), \tag{3.23}$$

$$\hat{\nabla}_i \hat{E}^i = 0, \tag{3.24}$$

where  $\hat{\nabla}$  is the covariant derivative of the metric  $\hat{h}$ . Thus the hypotheses of theorem 3.3 are satisfied. Since the mass of  $\hat{h}$  and the charge of  $\hat{E}$  coincide, theorem 3.3 shows that  $\hat{h}$  is, locally, the space part of the Majumdar–Papapetrou† metric. This shows in particular that  $\hat{h}$ , and hence also  $h$ , are conformally flat, so that the Cotton tensor  $B_{ijk}$  of  $h$  satisfies

$$B_{ijk} \equiv 0. \tag{3.25}$$

Equation (3.10) implies  $d\hat{E} = 0$  and from equation (3.22) we have

$$0 = d\hat{E} = 2 \frac{d\phi \wedge dV}{V^2}. \tag{3.26}$$

It follows that  $d\phi$  is parallel to  $dV$  wherever  $dV$  does not vanish. Standard results about solutions of elliptic equations show that

$$dV = -\frac{m \, dr}{r^2} + O(r^{-3}), \tag{3.27}$$

so that  $dV$  does not vanish for  $r \geq R$ , for  $R$  large enough. Increasing  $R$  if necessary it follows from equation (3.19) and from the maximum principle that for  $r \geq R$  the level sets of  $V$  will be embedded spheres. One also finds that there exists  $0 \leq V_- < 1$  such that for  $c \in [V_-, 1)$  the level sets  $\{V = c\}$  are smooth embedded spheres. Let

$$\begin{aligned} \hat{I} &= \{c \mid c \text{ is a non-critical value of } V\}, \\ \hat{\mathcal{U}} &= \{p \mid V(p) \in \hat{I}\} = \cup_{c \in \hat{I}} V^{-1}(c), \end{aligned}$$

and define  $\mathcal{U}$  to be that connected component of  $\hat{\mathcal{U}}$  that contains  $\mathbb{R}^3 \setminus B(0, R)$ . (Recall that  $c$  is non-critical if  $dV$  is nowhere vanishing on the level set  $V = c$ .) Similarly define  $I \subset (0, 1)$  to be that connected component of  $\hat{I} \setminus \{0\}$  that contains  $(V_-, 1)$ ; clearly,

$$\mathcal{U} = \{p \mid V(p) \in I\} = \cup_{c \in I} V^{-1}(c).$$

Compactness of the level sets of  $V$  implies that  $\mathcal{U}$  is diffeomorphic to  $I \times S^2$ , and that on  $\mathcal{U}$  the function  $V$  can be used as a coordinate. Further, we can introduce a finite number of coordinate patches with coordinates  $x^A$ ,  $A = 1, 2$ , on  $S^2$  so that on  $\mathcal{U}$  the metric takes the form

$$h = W^{-2} dV^2 + \gamma_{AB} dx^A dx^B. \tag{3.28}$$

Equation (3.26) shows that

$$\phi = \phi(V)$$

on  $\mathcal{U}$ . This allows one to write equations (3.5) and (3.7) in the coordinate system (3.28) as

$$\frac{\partial \phi}{\partial V} = \frac{V}{W \sqrt{\det \gamma_{AB}}} \frac{\partial}{\partial V} \left( W \sqrt{\det \gamma_{AB}} \frac{\partial \phi}{\partial V} \right), \tag{3.29}$$

$$\left( \frac{\partial \phi}{\partial V} \right)^2 = \frac{V}{W \sqrt{\det \gamma_{AB}}} \frac{\partial}{\partial V} \left( W \sqrt{\det \gamma_{AB}} \right). \tag{3.30}$$

It follows that

$$V \frac{\partial^2 \phi}{\partial V^2} = \frac{\partial \phi}{\partial V} - \left( \frac{\partial \phi}{\partial V} \right)^3.$$

† It might be worthwhile to point out that it is not known at this stage that  $\hat{h}$  is the space part of a standard Majumdar–Papapetrou metric, but this information is not needed in the argument.

Integrating this equation one finds

$$\frac{\partial\phi}{\partial V} = \frac{AV}{\sqrt{1+A^2V^2}}, \quad (3.31)$$

where  $A$  is an integration constant. From equation (3.27), from  $m > 0$  and from  $d\phi = -Q dr/r^2 + O(r^{-3})$  one obtains

$$\lim_{V \rightarrow 1} \frac{\partial\phi}{\partial V} = -\frac{Q}{m},$$

which determines  $A$ .

Suppose first that  $Q = 0$ , then

$$d\phi \equiv 0$$

on  $\mathcal{U}$  by equation (3.31). Equation (3.5) and the unique continuation theorem of Aronszajn [1] show that  $\phi$  is constant throughout  $\Sigma$ , so that the initial data set is vacuum. In this case the spacetime metric is the Schwarzschild metric in a neighbourhood of  $\Sigma$  by theorem 1.1 of [10].

It remains to consider the case  $Q \neq 0$ . Integrating equation (3.31) and using  $\lim_{V \rightarrow 1} \phi = 0$  we obtain

$$\phi = \frac{m - \sqrt{m^2 + Q^2(V^2 - 1)}}{Q}. \quad (3.32)$$

According to Heusler [20] (equation (9.58); cf also [25]) this implies

$$\frac{V^4}{8W^4} B_{ijk} B^{ijk} = \left( \frac{m^2 - Q^2}{m^2 + Q^2(V^2 - 1)} \right)^2 (|\lambda|_\gamma^2 + \frac{1}{2} |\mathcal{D}W|_\gamma^2).$$

Here  $|\cdot|_\gamma$  denotes the norm with respect to the metric  $\gamma = \gamma_{AB} dx^A dx^B$ ,  $\mathcal{D}W$  is the gradient of the restriction of the function  $W$  (defined in (3.28)) to the level sets of  $V$ , and  $\lambda \equiv \lambda_{AB} dx^A dx^B$  is the trace-free part of the extrinsic curvature tensor of the level sets of  $V$ —in the coordinate system of (3.28)

$$\lambda_{AB} = W \left( \frac{\partial\gamma_{AB}}{\partial V} - \frac{1}{2} \gamma^{CD} \frac{\partial\gamma_{CD}}{\partial V} \gamma_{AB} \right). \quad (3.33)$$

Equation (3.25) implies that  $\partial\gamma_{AB}/\partial V$  is pure trace, and that  $W = W(V)$ . This latter property and equation (3.30) show that  $\det \gamma_{AB}$  is a product of a function of  $V$  with a function of the remaining coordinates. From the asymptotic behaviour of the metric it then follows that

$$h = W(V)^{-2} dV^2 + H(V) d\Omega^2, \quad (3.34)$$

for some function  $H(V)$ , where  $d\Omega^2$  is the standard round metric on  $S^2$ . A straightforward integration of equations (3.30) and (3.9) using (3.32) shows that the metric on  $\mathcal{U}$  is the space part of the Reissner–Nordström metric. In other words,  $h$  is on  $\mathcal{U}$  the pull back by a suitable diffeomorphism  $\psi$  of the space part  $h^{\text{RN}}$  of the Reissner–Nordström metric.

To finish the proof<sup>†</sup>, we claim that  $I$  is open in  $(0, 1)$ , which can be seen as follows. Let  $p \in \mathcal{U}$ , we thus have  $dV(q) \neq 0$  for all  $q$  such that  $V(p) = V(q)$ . By equation (3.34)  $|dV|_h = W$  is constant on the level set  $V^{-1}(V(p))$  of  $V$  through  $p$  so that

$$\inf_{V^{-1}(V(p))} |dV|_h > 0,$$

which easily implies that all nearby level sets are non-critical.

<sup>†</sup> We note that it is usual at this stage to invoke analyticity to conclude the proof. Because analytic extensions of manifolds are not unique this is not sufficient without further justification. The argument we present here avoids this problem.

To see that  $I$  is closed in  $(0, 1)$ , recall that, using obvious notation, we have  $h = \psi^* h^{\text{RN}}$  and  $V = V^{\text{RN}} \circ \psi$  on  $\mathcal{U}$ . Let  $s_i \in I$  be any sequence converging to  $s \in (0, 1)$ , thus  $s_i = V(p_i)$  for some  $p_i \in \mathcal{U}$ . By the interior compactness of  $\Sigma$ , passing to a subsequence if necessary, there exists  $p \in \Sigma$  such that  $p_i \rightarrow p$ , with  $V(p) = s > 0$ . Set

$$C = \inf |dV^{\text{RN}}|_{h^{\text{RN}}},$$

where the infimum is taken over those points  $q$  in  $M^{\text{RN}}$  for which  $V^{\text{RN}}(q) > V(p)/2$ . We have  $V(p_i) = s_i > V(p)/2$  for  $i$  large enough, so that  $V^{\text{RN}}(\psi(p_i)) = V(p_i) > V(p)/2$ . It follows that

$$|dV|_h(p) = \lim_{i \rightarrow \infty} |dV|_h(p_i) = \lim_{i \rightarrow \infty} |dV^{\text{RN}}|_{h^{\text{RN}}}(\psi(p_i)) > C,$$

so that  $dV(p) \neq 0$ . Now  $|dV|_h$  is constant on those level sets of  $V$  which are in  $I$ , and by continuity it is also constant on those level sets of  $V$  which are in  $\bar{I}$ , the closure of  $I$  in  $(0, 1)$ . Hence  $|dV|_h$  is non-vanishing on the level set  $\{V = s\}$ , thus  $s \in I$ .

We have thus shown that  $I$  is open and closed, and connectedness of  $\Sigma$  implies  $\mathcal{U} = \Sigma$ . Thus the manifold  $\mathbb{R} \times \Sigma$  with the metric  $-V^2 dt^2 + h$  is isometrically diffeomorphic to the Reissner–Nordström spacetime.

Consider any neighbourhood  $\mathcal{V}$  of  $\Sigma$  diffeomorphic to an open interval times  $\Sigma$ ; the set  $\mathcal{V}$  is simply connected by the simple-connectedness of  $\Sigma$ . Let  $\alpha$  be the 1-form

$$\alpha = \frac{X_\mu dx^\mu}{X_\nu X^\nu}.$$

Equation (2.1) shows that  $\alpha$  is closed, and simple-connectedness of  $\mathcal{V}$  implies existence of a function  $t \in C^\infty(\mathcal{V})$  such that  $\alpha = dt$ . As in the proof of lemma 5.1 of [10] there exists a function  $f: \Sigma \rightarrow \mathbb{R}$  such that

$$t = s + f. \tag{3.35}$$

Here  $s$  denotes the coordinate along the (perhaps only locally defined) orbits of the Killing vector field on  $\mathcal{V}$ . Passing to a subset of  $\mathcal{V}$  if necessary we may assume that every orbit of  $X$  in  $\mathcal{V}$  intersects  $\Sigma$  precisely once. We can then extend  $f$  to a function on  $\mathcal{V}$  by requiring that  $X(f) = 0$ . As the metric  $-V^2 dt^2 + h$  has already been shown to be the Reissner–Nordström metric, equation (3.35) now provides the required embedding of  $\mathcal{V}$  into an open subset of the Reissner–Nordström spacetime.  $\square$

In order to show that theorem 1.1 is a special case of theorem 3.6 we need the following result.

**Lemma 3.7.** *Let  $S_a$ ,  $a = 1, 2$  be connected components of  $\partial\Sigma$  such that the horizon potentials  $\phi_a = \phi|_{S_a}$  satisfy*

$$\phi_1 = \inf_{\bar{\Sigma}} \phi < 0, \quad \phi_2 = \sup_{\bar{\Sigma}} \phi > 0. \tag{3.36}$$

*Then the charges  $Q_a$  of the  $S_a$ 's are non-vanishing and have opposite signs.*

The result is obtained by standard integration by parts arguments. However, some care must be taken in our context because the degenerate components of the boundary  $\partial\Sigma$  lie at infinite  $h$ -distance, and because  $V$  tends to zero there.

**Proof.** Recall that the charges of the  $S_a$ 's can be defined by the equations

$$Q_a = - \lim_{i \rightarrow \infty} \int_{S_{a,i}} V^{-1} \nabla^i \phi dS_i, \tag{3.37}$$

where the  $S_{a,i}$  are any family of connected smooth hypersurfaces converging in an appropriate sense to the  $S_a$ 's as  $i$  tends to infinity. For definiteness:

- If  $S_a$  is degenerate we take the  $S_{a,i}$ 's to be the sets  $x = 1/i$ , where  $x$  is the coordinate of the proof of proposition 3.2 of [10], and we assume that  $x$  has been rescaled so that its range covers the interval  $[0, 1]$ ; we set

$$\Omega_a = \{x < 1\}.$$

- If  $S_a$  is non-degenerate we take the  $S_{a,i}$ 's to be the sets  $w = 1/i$ , where  $w$  is the coordinate of the proof of proposition 3.3 of [10], and we assume that  $w$  has been rescaled so that its range covers the interval  $[0, 1]$ ; we set

$$\Omega_a = \{w < 1\}.$$

The integrals on the right-hand side of equation (3.37) are  $i$  independent by equation (3.5) and the divergence theorem,

$$\int_{S_{a,i}} V^{-1} \nabla^i \phi \, dS_i - \int_{S_{a,j}} V^{-1} \nabla^i \phi \, dS_i = \int_{\mathcal{V}_{i,j}} \nabla_i (V^{-1} \nabla^i \phi) \, d\mu_h = 0.$$

Here  $\mathcal{V}_{i,j}$  is the volume with a boundary consisting of  $S_{a,i}$  and  $S_{a,j}$ . Hence the limit in (3.37) exists.

Let

$$\phi_- = \inf_{S_{1,1}} \phi, \quad \phi_+ = \sup_{S_{2,1}} \phi.$$

From equation (3.36) and the maximum principle we have  $\phi_1 < \phi_-$  and  $\phi_+ < \phi_2$ . Let  $c$  be a non-critical value of  $\phi$  satisfying  $\phi_1 < c < \phi_-$ , then the level set  $\phi^{-1}\{c\} \cap \Omega_1$  is a smooth compact submanifold of  $\Omega_1$ ; recall that the set of non-critical values of  $\phi$  is dense by Sard's theorem (cf, for example, [15]). Applying the divergence theorem on a set bounded by  $\phi^{-1}\{d\} \cap \Omega_1$  (with a non-critical  $d$  satisfying  $\phi_1 < d < \phi_-$ ) and by  $S_{1,i}$  for an  $i$  large enough we obtain

$$\int_{\phi^{-1}\{d\} \cap \Omega_1} V^{-1} \nabla^i \phi \, dS_i = -Q_1. \quad (3.38)$$

Let  $c$  and  $d$  be any non-critical values of  $\phi$  satisfying  $\phi_1 < c < d < \phi_-$ , thus  $\mathcal{W}_{cd} \equiv \{c \leq \phi \leq d\} \cap \Omega_1$  is a smooth compact submanifold of  $\Omega_1$  with boundary  $(\phi^{-1}\{c\} \cup \phi^{-1}\{d\}) \cap \Omega_1$ . By the maximum principle and the boundary point lemma (lemma 3.4 of [14]) we have

$$h(\nabla \phi, n) > 0$$

on  $\phi^{-1}\{d\} \cap \Omega_1$ , where  $n$  is that unit normal to  $\phi^{-1}\{d\} \cap \Omega_1$  which points outwards from  $\mathcal{W}_{cd}$ , hence

$$Q_1 = - \int_{\phi^{-1}\{d\} \cap \Omega_1} V^{-1} \nabla^i \phi \, dS_i < 0.$$

The inequality  $Q_2 > 0$  follows by changing  $\phi$  to  $-\phi$  in the argument above.  $\square$

We can now pass to the

**Proof of theorem 1.1.** If  $\partial \Sigma$  is connected the hypotheses of theorem 3.6 are obviously satisfied, and the result follows. Suppose, thus, that  $\partial \Sigma$  is not connected. Changing  $\phi$  to  $-\phi$  if necessary we will be able to satisfy (3.20) unless there exists a connected component  $S_1$  of  $\partial \Sigma$  such that  $\phi_1 = -1$  and a connected component  $S_2$  of  $\partial \Sigma$  such that  $\phi_2 = 1$ . By proposition 3.4  $S_1$  and  $S_2$  have to be degenerate, and by lemma 3.7 the charges of  $S_1$  and  $S_2$  have opposite signs. This is, however, not allowed by the hypotheses of theorem 1.1, and the result follows by theorem 3.6.  $\square$

We finally note that the Reissner–Nordström case of corollary 1.2 is proved by a repetition of the arguments of the proof of corollary 1.2 in [10]. The Majumdar–Papapetrou case is proved by a repetition of the arguments of the proof of corollary 1.2 in [10] together with the arguments presented in the first paragraph of the proof of theorem 3.6.

## Acknowledgment

The author acknowledges useful discussions with W Simon.

## References

- [1] Aronszajn N 1957 A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order *J. Math.* **XXXVI** 235–49
- [2] Ashtekar A and Magnon-Ashtekar A 1979 On conserved quantities in general relativity *J. Math. Phys.* **20** 793–800
- [3] Bartnik R and Chruściel P T 1998 in preparation
- [4] Beig R 1978 Arnowitt–Deser–Misner energy and  $g_{00}$  *Phys. Lett. A* **69** 153–5
- [5] Beig R and Chruściel P T 1996 Killing vectors in asymptotically flat space-times: I. Asymptotically translational Killing vectors and the rigid positive energy theorem *J. Math. Phys.* **37** 1939–61
- [6] Carter B 1969 Killing horizons and orthogonally transitive groups in space-time *J. Math. Phys.* **10** 70–81
- [7] Chruściel P T 1986 A remark on the positive energy theorem *Class. Quantum Grav.* **33** L115–21
- [8] Chruściel P T 1994 ‘No hair’ theorems—folklore, conjectures, results *Differential Geometry and Mathematical Physics (Cont. Math. vol 170)* ed J Beem and K L Duggal (Providence, RI: American Mathematical Society) pp 23–49
- [9] Chruściel P T 1996 Uniqueness of black holes revisited *Proc. Journées Relativistes 96* ed N Straumann, Ph Jetzer and G Lavrelashvili (*Helv. Phys. Acta* **69** 529–52)
- [10] Chruściel P T 1999 The classification of static and vacuum spacetimes containing an asymptotically flat spacelike hypersurface with compact interior *Class. Quantum Grav.* **16** 661–87
- [11] Chruściel P T and Nadirashvili N S 1995 All electrovacuum Majumdar–Papapetrou spacetimes with non-singular black holes *Class. Quantum Grav.* **12** L17–23
- [12] Chruściel P T and Wald R M 1994 On the topology of stationary black holes *Class. Quantum Grav.* **11** L147–52
- [13] Gibbons G W and Hull C M 1982 A Bogomolny bound for general relativity and solitons in  $N = 2$  supergravity *Phys. Lett. B* **109** 190–4
- [14] Gilbarg D and Trudinger N S 1983 *Elliptic Partial Differential Equations of Second Order* (Berlin: Springer)
- [15] Guillemin V and Pollack A 1974 *Differential Topology* (Englewood Cliffs, NJ: Prentice-Hall)
- [16] Hartle J B and Hawking S W 1972 Solutions of the Einstein–Maxwell equations with many black holes *Commun. Math. Phys.* **26** 87–101
- [17] Hawking S W and Ellis G F R 1973 *The Large Scale Structure of Space-Time* (Cambridge: Cambridge University Press)
- [18] Herzlich M 1998 The positive mass theorem for black holes revisited *J. Geom. Phys.* **26** 97–111
- [19] Heusler M 1994 On the uniqueness of the Reissner–Nordström solution with electric and magnetic charge *Class. Quantum Grav.* **11** L49–53
- [20] Heusler M 1996 *Black Hole Uniqueness Theorems* (Cambridge: Cambridge University Press)
- [21] Heusler M 1997 On the uniqueness of the Papapetrou–Majumdar metric *Class. Quantum Grav.* **14** L129–34
- [22] Israel W 1968 Event horizons in static electrovac space-times *Commun. Math. Phys.* **8** 245–60
- [23] Majumdar S D 1947 A class of exact solutions of Einstein’s field equations *Phys. Rev.* **72** 390–8
- [24] Masood-ul-Alam A K M 1992 Uniqueness proof of static charged black holes revisited *Class. Quantum Grav.* **9** L53–5
- [25] Müller zum Hagen H, Robinson D C and Seifert H J 1974 Black holes in static electrovac space-times *Gen. Rel. Grav.* **5** 61–72
- [26] Papapetrou A 1947 A static solution of the equations of the gravitational field for an arbitrary charge distribution *Proc. R. Irish Acad A* **51** 191–204
- [27] Ruback P 1988 A new uniqueness theorem for charged black holes *Class. Quantum Grav.* **5** L155–9
- [28] Simon W 1992 Radiative Einstein–Maxwell spacetimes and ‘no-hair’ theorems *Class. Quantum Grav.* **9** 241–56
- [29] Sudarsky D and Wald R M 1993 Extrema of mass, stationarity and staticity, and solutions to the Einstein–Yang–Mills equations *Phys. Rev. D* **46** 1453–74

- [30] Tod K P 1983 All metrics admitting super-covariantly constant spinors *Phys. Lett. B* **121** 241–4
- [31] Vishveshwara C V 1968 Generalization of the ‘Schwarzschild surface’ to arbitrary static and stationary metrics  
*J. Math. Phys.* **9** 1319–22