

# Fully Complementary Higher Dimensional Partitions

Florian Schreier-Aigner  
University of Vienna

90th SLC, Bad Boll

# Partitions

---

A **partition**  $\lambda = (\lambda_1, \lambda_2, \dots,)$  is a weakly decreasing sequence of non-negative integers with all but finitely many entries equal to 0. We define the size  $|\lambda| = \lambda_1 + \lambda_2 + \dots$ .

$$(4, 2, 1, 0, \dots)$$

# Partitions

A **partition**  $\lambda = (\lambda_1, \lambda_2, \dots)$  is a weakly decreasing sequence of non-negative integers with all but finitely many entries equal to 0. We define the size  $|\lambda| = \lambda_1 + \lambda_2 + \dots$ .

$$(4, 2, 1, 0, \dots) \Leftrightarrow \begin{array}{|c|c|c|c|} \hline 1, 1 & 1, 2 & 1, 3 & 1, 4 \\ \hline 2, 1 & 2, 2 & & \\ \hline 3, 1 & & & \\ \hline \end{array}$$

A **Young diagram**  $\lambda$  is a finite subset of  $\mathbb{N}_{>0}^2$  such that  $(x_1, x_2) \in \lambda$  implies  $(y_1, y_2) \in \lambda$  for  $1 \leq y_i \leq x_i$  for  $1 \leq i \leq 2$ .

# Generating functions I

## Theorem

*The generating function for partitions is*

$$\sum_{\lambda} q^{|\lambda|} = \prod_{i \geq 1} \frac{1}{1 - q^i}.$$

# Generating functions I

## Theorem

The generating function for partitions is

$$\sum_{\lambda} q^{|\lambda|} = \prod_{i \geq 1} \frac{1}{1 - q^i}.$$

A Young diagram  $\lambda$  is **contained** in an  $(a, b)$ -box if  $\lambda \subseteq [a] \times [b]$ .

## Theorem

The generating function of Young diagrams inside an  $(a, b)$ -box is

$$\sum_{\lambda} q^{|\lambda|} = \left[ \begin{matrix} a + b \\ a \end{matrix} \right]_q = \prod_{i=1}^a \prod_{j=1}^b \frac{1 - q^{i+j}}{1 - q^{i+j-1}}.$$

# Plane partitions


---

A **plane partition**  $\pi$  is an array  $(\pi_{i,j})$  of non-negative integers and finite support, which is weakly decreasing along rows and columns.

$$\begin{array}{cccc} 3 & 2 & 2 & 1 \\ 2 & 2 & 1 & \\ 1 & & & \end{array}$$

# Plane partitions

A **plane partition**  $\pi$  is an array  $(\pi_{i,j})$  of non-negative integers and finite support, which is weakly decreasing along rows and columns.

$$\begin{array}{cccc} 3 & 2 & 2 & 1 \\ 2 & 2 & 1 & \\ 1 & & & \end{array}$$
$$\Leftrightarrow$$


A **2-dimensional Young diagram**  $\lambda$  is a finite subset of  $\mathbb{N}_{>0}^3$  such that  $(x_1, x_2, x_3) \in \lambda$  implies  $(y_1, y_2, y_3) \in \lambda$  for  $1 \leq y_i \leq x_i$  for  $1 \leq i \leq 3$ .

## Generating functions II

### Theorem (MacMahon)

*The generating function for plane partitions is*

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^i}.$$



## Generating functions II

### Theorem (MacMahon)

*The generating function for plane partitions is*

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^i}.$$

A 2-dimensional Young diagram  $\lambda$  is **contained** in an  $(a, b, c)$ -box if  $\lambda \subseteq [a] \times [b] \times [c]$ .

### Theorem (MacMahon)

*The generating function for 2-dimensional Young diagrams inside an  $(a, b, c)$ -box is*

$$\sum_{\lambda} q^{|\lambda|} = \prod_{i=1}^a \prod_{j=1}^b \prod_{k=1}^c \frac{1 - q^{i+j+k-1}}{1 - q^{i+j+k-2}}.$$

# $d$ -dimensional partitions

A  **$d$ -dimensional partition**  $\pi$  is an array  $(\pi_{i_1, \dots, i_d})$  of non-negative integers and finite support, such that

$$\pi_{i_1, \dots, i_d} \geq \pi_{i_1, \dots, i_k+1, \dots, i_d},$$

for all  $i_1, \dots, i_d \in \mathbb{N}_{>0}$  and  $1 \leq k \leq d$ .

$$\begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array} \quad \begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array}$$

## $d$ -dimensional partitions

A  **$d$ -dimensional partition**  $\pi$  is an array  $(\pi_{i_1, \dots, i_d})$  of non-negative integers and finite support, such that

$$\pi_{i_1, \dots, i_d} \geq \pi_{i_1, \dots, i_k+1, \dots, i_d},$$

for all  $i_1, \dots, i_d \in \mathbb{N}_{>0}$  and  $1 \leq k \leq d$ .

$$\begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array} \quad \begin{array}{cc} 1 & 0 \\ 0 & 0 \end{array} \quad \Leftrightarrow \quad \{(1, 1, 1, 1), (1, 1, 2, 1)\}$$

A  **$d$ -dimensional Young diagram**  $\lambda$  is a finite subset of  $\mathbb{N}_{>0}^{d+1}$  such that  $\mathbf{x} \in \lambda$  implies  $\mathbf{y} \in \lambda$  for  $1 \leq y_i \leq x_i$  for  $1 \leq i \leq d+1$ .

# Generating functions III

## Conjecture (MacMahon)

*The generating function of  $d$ -dimensional partitions  $\pi$  is*

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^{\binom{d+i-2}{d-1}}}.$$

# Generating functions III

## Conjecture (MacMahon)

*The generating function of  $d$ -dimensional partitions  $\pi$  is*

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^{\binom{d+i-2}{d-1}}}.$$

This was disproved by Atkin, Bratley, Macdonald and McKay in 1967.

# Generating functions III

## Conjecture (MacMahon)

*The generating function of  $d$ -dimensional partitions  $\pi$  is*

$$\sum_{\pi} q^{|\pi|} = \prod_{i \geq 1} \frac{1}{(1 - q^i)^{\binom{d+i-2}{d-1}}}.$$



This was disproved by Atkin, Bratley, Macdonald and McKay in 1967.

## Theorem (Amanov–Yeliussizov, 2023)


*The generating function of  $d$ -dimensional partitions  $\pi$  with respect to two statistics  $\text{cor}$  and  $|\cdot|_{ch}$  is given by*

$$\sum_{\pi} t^{\text{cor}(\pi)} q^{|\pi|_{ch}} = \prod_{i \geq 1} (1 - tq^i)^{-\binom{i+d-2}{d-1}}.$$


# Symmetries of boxed plane partitions




reflection



rotation



complementation



# Self-complementary VS fully complementary



---

A  $2d$ -Young diagram  $\lambda$  inside an  $(a, b, c)$ -box is called **self-complementary** if it is equal to its complementation.






# Self-complementary VS fully complementary

A  $2d$ -Young diagram  $\lambda$  inside an  $(a, b, c)$ -box is called **self-complementary** if  $\lambda$ , and  $\lambda$  “placed” at the corner  $(a, b, c)$  fills the  $(a, b, c)$ -box without overlap.



# Self-complementary VS fully complementary

A  $2d$ -Young diagram  $\lambda$  inside an  $(a, b, c)$ -box is called **self-complementary** if  $\lambda$ , and  $\lambda$  “placed” at the corner  $(a, b, c)$  fills the  $(a, b, c)$ -box without overlap.



A  $2d$ -Young diagram  $\lambda$  inside an  $(a, b, c)$ -box is called **fully complementary** if  $\lambda$ , and  $\lambda$  “placed” at the corners  $(a, b, 1)$ ,  $(a, 1, c)$  and  $(1, b, c)$  fill the  $(a, b, c)$ -box without overlap.

# Fully Complementary in higher dimensions

Let  $\mathbf{n} = (n_1, \dots, n_{d+1})$  be a sequence of positive integers and  $I \subseteq [d+1] = \{1, 2, \dots, d+1\}$ . We define for  $\mathbf{x} = (x_1, \dots, x_{d+1})$

$$\rho_{I, \mathbf{n}}(\mathbf{x}) := \left( \begin{cases} x_i & i \notin I, \\ 2n_i + 1 - x_i & i \in I, \end{cases} \right)_{1 \leq i \leq d+1}.$$

A  $d$ -dimensional Young diagram  $\lambda$  is called **fully complementary** inside a  $(2n_1, \dots, 2n_{d+1})$ -box if

- for all even sized  $I \neq J \subseteq [d+1]$  holds  $\rho_{I, 2\mathbf{n}}(\lambda) \cap \rho_{J, 2\mathbf{n}}(\lambda) = \emptyset$ ,
- and  $\bigcup_{\substack{I \subseteq [d+1] \\ |I| \text{ even}}} \rho_{I, 2\mathbf{n}}(\lambda) = [2n_1] \times \dots \times [2n_{d+1}]$ .

## An example

For  $\mathbf{n} = (1, 1, 1, 1)$  the 3-dimensional Young diagram  $\lambda = \{(1, 1, 1, 1), (2, 1, 1, 1)\}$  is fully complementary inside the  $(2, 2, 2, 2)$ -box.

$$\rho_{\{1,2\}}(\lambda) = \{(2, 2, 1, 1), (1, 2, 1, 1)\},$$

$$\rho_{\{2,3\}}(\lambda) = \{(1, 2, 2, 1), (2, 2, 2, 1)\},$$

$$\rho_{\{1,3\}}(\lambda) = \{(2, 1, 2, 1), (1, 1, 2, 1)\},$$

$$\rho_{\{2,4\}}(\lambda) = \{(1, 2, 1, 2), (2, 2, 1, 2)\},$$

$$\rho_{\{1,4\}}(\lambda) = \{(2, 1, 1, 2), (1, 1, 1, 2)\},$$

$$\rho_{\{3,4\}}(\lambda) = \{(1, 1, 2, 2), (2, 1, 2, 2)\},$$

$$\rho_{\{1,2,3,4\}}(\lambda) = \{(2, 2, 2, 2), (1, 2, 2, 2)\}.$$

## An example

For  $\mathbf{n} = (1, 1, 1, 1)$  the 3-dimensional Young diagram  $\lambda = \{(1, 1, 1, 1), (2, 1, 1, 1)\}$  is fully complementary inside the  $(2, 2, 2, 2)$ -box.

$$\begin{aligned}\rho_{\{1,2\}}(\lambda) &= \{(2, 2, 1, 1), (1, 2, 1, 1)\}, & \rho_{\{2,3\}}(\lambda) &= \{(1, 2, 2, 1), (2, 2, 2, 1)\}, \\ \rho_{\{1,3\}}(\lambda) &= \{(2, 1, 2, 1), (1, 1, 2, 1)\}, & \rho_{\{2,4\}}(\lambda) &= \{(1, 2, 1, 2), (2, 2, 1, 2)\}, \\ \rho_{\{1,4\}}(\lambda) &= \{(2, 1, 1, 2), (1, 1, 1, 2)\}, & \rho_{\{3,4\}}(\lambda) &= \{(1, 1, 2, 2), (2, 1, 2, 2)\}, \\ \rho_{\{1,2,3,4\}}(\lambda) &= \{(2, 2, 2, 2), (1, 2, 2, 2)\}.\end{aligned}$$

There are three further Young diagrams which are fully complementary inside  $(2, 2, 2, 2)$ :

$$\{(1, 1, 1, 1), (1, 2, 1, 1)\}$$

$$\{(1, 1, 1, 1), (1, 1, 2, 1)\}$$

$$\{(1, 1, 1, 1), (1, 1, 1, 2)\}$$

## Generating functions IV

We call a  $d$ -dimensional partition  $\pi$  **fully complementary inside a  $(2n_1, \dots, 2n_{d+1})$ -box** if its Young diagram is fully complementary in this box.

### Theorem (SA, 2023)

Let  $\mathbf{x} = (x_1, \dots, x_{d+1})$ ,  $\mathbf{n} = (n_1, \dots, n_{d+1}) \in \mathbb{N}_{>0}^{d+1}$  and denote by  $\text{FCP}(\mathbf{n})$  the set of fully complementary partitions inside a  $(2n_1, \dots, 2n_{d+1})$ -box. Then

$$\sum_{\mathbf{n} \in \mathbb{N}_{>0}^{d+1}} |\text{FCP}(\mathbf{n})| \mathbf{x}^{\mathbf{n}} = \frac{\left( d + 1 - \sum_{i=1}^{d+1} x_i \right) \prod_{i=1}^{d+1} x_i}{\left( 1 - \sum_{i=1}^{d+1} x_i \right) \prod_{i=1}^{d+1} (1 - x_i)}.$$

## Stretching maps

For  $1 \leq k \leq d$  define the map  $\varphi_k : \text{FCP}(\mathbf{n}) \rightarrow \text{FCP}(\mathbf{n} + e_k)$

$$\varphi_k(\pi)_{i_1, \dots, i_d} = \begin{cases} \pi_{i_1, \dots, i_d} & i_k \leq n_k, \\ n_{d+1} & i_k \in \{n_k + 1, n_k + 2\} \\ & \text{and } i_j \leq n_j \text{ for all } 1 \leq j \neq k \leq d, \\ \pi_{i_1, \dots, i_k - 2, \dots, i_d} & i_k > n_k + 2, \\ 0 & \text{otherwise,} \end{cases}$$

and the map  $\varphi_{d+1} : \text{FCP}(\mathbf{n}) \rightarrow \text{FCP}(\mathbf{n} + e_{d+1})$  as

$$\varphi_{d+1}(\pi)_{i_1, \dots, i_d} = \begin{cases} \pi_{i_1, \dots, i_d} + 2 & i_j \leq n_j \text{ for all } 1 \leq j \leq d, \\ \pi_{i_1, \dots, i_d} & \text{otherwise.} \end{cases}$$

# A recursive structure

## Proposition

Let  $\mathbf{n} = (n_1, \dots, n_{d+1})$  be a sequence of positive integers. Then  $\text{FCP}(\mathbf{n})$  is equal to the disjoint union

$$\text{FCP}(\mathbf{n}) = \dot{\bigcup}_{1 \leq k \leq d+1} \varphi_k(\text{FCP}(\mathbf{n} - \mathbf{e}_k)).$$

Note, that if exactly one  $n_i = 0$ , we have to define  $\text{FCP}(\mathbf{n})$  to consist of “the empty array” and extend the definitions of the  $\varphi_k$  appropriately.



# Symmetry classes of FCPs I

---

There are many interesting symmetry classes of plane partitions. Is the same true for fully complementary plane partitions?

# Symmetry classes of FCPs I

---

There are many interesting symmetry classes of plane partitions. Is the same true for fully complementary plane partitions?

symmetric	⇒	quasi symmetric
cyclically symmetric	⇒	?
self-complementary	⇒	self-complementary
transpose-complementary	⇒	quasi transpose-complementary

# Symmetry classes of FCPs II

## Proposition

- 1 Denote by  $QS(a, c)$  the set of quasi symmetric fully complementary plane partitions (FCPP) inside an  $(a, a, c)$ -box, then holds

$$\sum_{a, c \geq 0} |QS(a, c)| x^a y^c = \frac{x + y - 2x^2 - xy}{(1-x)(1-2x-y)}.$$

- 2 The number of self-complementary FCPPs inside an  $(2a, 2b, 2c)$ -box is  $\binom{a+b}{a}$ .
- 3 The number of quasi transpose-complementary FCPPs inside an  $(2a, 2a, 2c)$ -box is  $2^a$ .

## Quasi transpose complementary PPs

A plane partition  $\pi$  inside an  $(a, a, c)$ -box is called **quasi transpose complementary** if

$$\pi_{i,j} + \pi_{a+1-j, a+1-i} = c,$$

holds for all  $1 \leq i, j \leq n$  with  $i + j \neq a + 1$ .

### Theorem

*The number of quasi transpose-symmetric plane partitions inside an  $(a, a, c)$ -box is equal to the number of symmetric plane partitions inside an  $(a, a, c)$ -box.*


As a side product we stumble upon the relation

$$2^{n-1} \text{TCPP}(n, n, 2c) = \text{SPP}(n-1, n-1, 2c+1).$$

I have also some conjectures for other quasi symmetry classes.


# Proof idea

6	6	6	5	4
6	5	3	3	1
6	5	3	3	0
6	4	1	1	0
1	0	0	0	0



# Proof idea

6	6	6	5	4
6	5	3	3	1
6	5	3	3	0
6	4	1	1	0
1	0	0	0	0





# Proof idea

6	6	6	5	4
6	5	3	3	1
6	5	3	3	0
6	4	1	1	0
5	0	0	0	0

