

Ivrea-Verbano zone (Italy, Switzerland)

„Petrological“ reference list (\geq 2000)

- Abart, R., Schmud, R., & Harlov, D. (2001). Metasomatism coronas around hornblendite xenoliths in granulite facies marble, Ivrea zone, N Italy; I, constraints on component mobility. Contributions to Mineralogy and Petrology, 141(4), 473-493. doi:<http://dx.doi.org/10.1007/s004100100255>
- Abart, R., & Sperb, R. (2001). Metasomatic coronas around hornblendite xenoliths in granulite facies marble, Ivrea zone, N Italy; II, oxygen isotope patterns. Contributions to Mineralogy and Petrology, 141(4), 494-504. doi:<http://dx.doi.org/10.1007/s004100100256>
- Albertini, C. (2003). Il diopside del Monte Cervandone (Alpe Devero, Verbano-Cusio-Ossola). Rivista Mineralogica Italiana, 27(3), 126-133.
- Ando, S., Morton, A., & Garzanti, E. (2013). Metamorphic grade of source rocks revealed by chemical fingerprints of detrital amphibole and garnet. Special Publication - Geological Society of London, 386(1), 351-371. doi:<http://dx.doi.org/10.1144/SP386.5>
- Badanina, I. Y., Malitch, K. N., & Belousova, E. A. (2013). U-Pb and Hf isotope characteristics of zircon from chromitites at Finero. Mineralogical Magazine, 77(5), 639. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.2>
- Barberini, V., Burlini, L., & Zappone, A. (2007). Elastic properties, fabric and seismic anisotropy of amphibolites and their contribution to the lower crust reflectivity. Tectonophysics, 445(3-4), 227-244. doi:<http://dx.doi.org/10.1016/j.tecto.2007.08.017>
- Berger, A., Mercolli, I., Kapferer, N., & Fügenschuh, B. (2012). Single and double exhumation of fault blocks in the internal Sesia-Lanzo zone and the Ivrea-Verbano zone (Biella, Italy). International Journal of Earth Sciences = Geologische Rundschau, 101(7), 1877-1894. doi:<http://dx.doi.org/10.1007/s00531-012-0755-6>
- Bethoux, N., Sue, C., Paul, A., Virieux, J., Frechet, J., Thouvenot, F., & Cattaneo, M. (2007). Local tomography and focal mechanisms in the south-western Alps; comparison of methods and tectonic implications. Tectonophysics, 432(1-4), 1-19. doi:<http://dx.doi.org/10.1016/j.tecto.2006.10.004>
- Boriani, A., & Giobbi, E. (2004). Does the basement of western southern Alps display a tilted section through the continental crust? A review and discussion. Periodico Di Mineralogia, 73(2), 5-22.
- Brown, M. (2003). Metamorphism. GeoTimes, 48(7), 16-17.
- Caironi, V., Colombo, A., & Tunisi, A. (2004). Geochemical approach to characterization and source identification of the protoliths of metasedimentary rocks; an example from the Southern Alps. Periodico Di Mineralogia, 73(2), 109-118.
- Campostrini, I. (2001). Le miniere di Brosso. Monografie - Museo Regionale Di Scienze Naturali, 29, 185p.
- Cernok, A., Blaha, S., & Klötzli, U. (2010). Mineral chemistry and petrology of monazite and xenotime in a prograde metamorphic sequence in the kinzigite formation of the Ivrea zone, northern Italy. Geophysical Research Abstracts, 12, EGU2010-12553.
- Cesare, B., Acosta-Vigil, A., Ferrero, S., & Bartoli, O. (2011). Melt inclusions in migmatites and granulites. Journal of the Virtual Explorer, 38, 21. doi:<http://dx.doi.org/10.3809/jvirtex.2011.00268>
- Cesare, B., Ferrero, S., Bartoli, O., Acosta-Vigil, A., Turina, A., Poli, S., & Ewing, T. (2012). Nanogranites in anatetic metapelites; reading the message in the bottle. International Geological Congress, Abstracts = Congres Geologique International, Resumes, 34, 1892.
- Cesare, B., Ferrero, S., Bartoli, O., Acosta-Vigil, A., Turina, A., Poli, S., . . . Bodnar, R. (2012). Nanogranites in anatetic metapelites; building up the database. Mineralogical Magazine, 76(6), 1555.
- Cocherie, A., Rossi, P., Fanning, C. M., & Deloule, E. (2004). Devonian to jurassic thermochronology of the south European lower crust traced through ion-probe U-Pb geochronology of zircons from lower crust kinzigite. International Geological Congress, Abstracts = Congres Geologique International, Resumes, 32, Part 2, 1467.
- Correia, C. T., Sinigoi, S., Girardi, V. A. V., Mazzucchelli, M., Tassinari, C. C. G., & Giovanardi, T. (2012). The growth of large mafic intrusions; comparing Niquelandia and Ivrea igneous complexes. Lithos (Oslo), 155, 167-182. doi:<http://dx.doi.org/10.1016/j.lithos.2012.08.024>
- De Giusti, F., Dal Piaz, G. V., Massironi, M., & Schiavo, A. (2004). Carta geotonica della valle d'Aosta. Memorie Di Scienze Geologiche, 55, 129-149, 1 sheet.

- Denyszyn, S. W., Fiorentini, M. L., & Davis, A. M. (2016). A bigger tent for CAMP? geochronology and geochemistry of mineralized lower-crustal intrusions in NW Italy. V.M. Goldschmidt Conference - Program and Abstracts, 26, 648.
- Diehl, T., Husen, S., Kissling, E., & Deichmann, N. (2009). High-resolution 3-D P-wave model of the alpine crust. *Geophysical Journal International*, 179(2), 1133-1147. doi:<http://dx.doi.org/10.1111/j.1365-246X.2009.04331.x>
- Economos, R. C., Quick, J. E., Sinigoi, S., & de Silva, ,S.L. (2013). Zircon oxygen isotopes reveal Ivrea-Verbano zone source characteristics of the Sesia Valley caldera. American Geophysical Union Fall Meeting, 2013, Abstract V53A-2767.
- Enea, F. C., Taylor, J., Storey, C. D., Marschall, H., & Konrad-Schmolke, M. (2011). Testing the use of detrital rutile to investigate HP/UHP rocks. *Mineralogical Magazine*, 75(3), 811.
- Ewing, T. A., Rubatto, D., & Hermann, J. (2011). Insights into lower crustal evolution from Hf isotope and zr thermometry data for rutile. *Mineralogical Magazine*, 75(3), 824.
- Ewing, T. A., Rubatto, D., & Hermann, J. (2010). Testing the behaviour of Hf isotopes in rutile in a high-temperature metamorphic environment. *Abstracts - Geological Society of Australia*, 98, 118.
- Ewing, T. A., Hermann, J., & Rubatto, D. (2013). The robustness of the Zr-in-rutile and Ti-in-zircon thermometers during high temperature metamorphism (Ivrea-Verbano Zone, Northern Italy). *Contributions to Mineralogy and Petrology*, 165(4), 757-779. doi:<http://dx.doi.org/10.1007/s00410-012-0834-5>
- Ewing, T. A., Rubatto, D., Beltrando, M., & Hermann, J. (2015). Constraints on the thermal evolution of the Adriatic margin during Jurassic continental breakup; U/Pb dating of rutile from the Ivrea-Verbano zone, Italy. *Contributions to Mineralogy and Petrology*, 169(4), Article 44. doi:<http://dx.doi.org/10.1007/s00410-015-1135-6>
- Ferrando, S., Bernoulli, D., & Compagnoni, R. (2004). The Canavese zone (internal western Alps); a distal margin of Adria. *Schweizerische Mineralogische Und Petrographische Mitteilungen = Bulletin Suisse De Mineralogie Et Petrographie*, 84(3), 237-256.
- Fiorentini, M. L., & Beresford, S. W. (2008). Role of volatiles and metasomatized subcontinental lithospheric mantle in the genesis of magmatic Ni-Cu-PGE mineralization; insights from in situ H, Li, B analyses of hydromagmatic phases from the Valmaggia ultramafic pipe, Ivrea-Verbano zone (NW Italy). *Terra Nova*, 20(5), 333-340. doi:<http://dx.doi.org/10.1111/j.1365-3121.2008.00825.x>
- Fiorentini, M. L., Grieco, G., Ferrario, A., & Tunisi, A. (2002). Petrological and metallogenic outlines of the Valmaggia ultramafic pipe (Ivrea zone), NW Alps, Italy. *Periodico Di Mineralogia*, 71(3), 219-239.
- Franz, L., & Romer, R. L. (2007). Caledonian high pressure metamorphism in the Strona-Ceneri zone (Southern Alps of Southern Switzerland and Northern Italy). *Swiss Journal of Geosciences*, 100(3), 457-467. doi:<http://dx.doi.org/10.1007/s00015-007-1232-2>
- Garde, A. A., Boriani, A., & Sorensen, E. V. (2015). Crustal modelling of the Ivrea-Verbano zone in Northern Italy re-examined; co-seismic cataclasis versus extensional shear zones and sideways rotation. *Tectonophysics*, 662, 291-311. doi:<http://dx.doi.org/10.1016/j.tecto.2015.04.003>
- Garuti, G., Bea, F., Zaccarini, F., & Montero, P. (2001). Age, geochemistry and petrogenesis of the ultramafic pipes in the Ivrea zone, NW Italy. *Journal of Petrology*, 42(2), 433-457. doi:<http://dx.doi.org/10.1093/petrology/42.2.433>
- Giovanardi, T., Zanetti, A., Mazzucchelli, M., Tiepolo, M., Wu, F. W., Langone, A., & Vannucci, R. (2013). U-pb and hf isotopes in zircons from mantle chromitites of the Finero peridotite (Ivrea Verbano zone). *Mineralogical Magazine*, 77(5), 1177. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.7>
- Grieco, G., Ferrario, A., von Quadt, A., Köppel, V., & Mathez, E. A. (2001). The zircon-bearing chromitites of the phlogopite peridotite of Finero (Ivrea zone, Southern Alps); evidence and geochronology of a metasomatized mantle slab. *Journal of Petrology*, 42(1), 89-101. doi:<http://dx.doi.org/10.1093/petrology/42.1.89>
- Grieco, G., Ferrario, A., & Mathez, E. A. (2004). The effect of metasomatism on the Cr-PGE mineralization in the Finero complex, Ivrea zone, southern Alps. *Ore Geology Reviews*, 24(3-4), 299-314. doi:<http://dx.doi.org/10.1016/j.oregeorev.2003.05.004>
- Gualtieri, L., Serretti, P., & Morelli, A. (2014). Finite-difference P wave travel time seismic tomography of the crust and uppermost mantle in the Italian region. *Geochemistry, Geophysics, Geosystems - G3*, 15(1), 69- 88. doi:<http://dx.doi.org/10.1002/2013GC004988>
- Günes, Z., Klötzli, U., & Sinigoi, S. (2008). Evidence for a ca. 1600 ma lower crust in the Ivrea-Verbano zone, Italy. *Geophysical Research Abstracts*, 10, EGU2008-A-07133.
- Günes, Z., Klötzli, U., & Sinigoi, S. (2007). Further evidence for a two stage magmatic underplating event in the Ivrea-Verbano zone, Italy. *Geochimica Et Cosmochimica Acta*, 71(15), 1.
- Halick, M. A., Selverstone, J., Sharp, Z. D., & Barnes, J. (2009). Chlorine isotopic heterogeneity in mantle peridotites from the Ivrea zone. *EOS, Transactions, American Geophysical Union*, 90(52), Abstract V31F-03.

- Handy, M. R., & Stuenitz, H. (2002). Strain localization by fracturing and reaction weakening; a mechanism for initiating exhumation of subcontinental mantle beneath rifted margins. Geological Society Special Publications, 200, 387-407.
- Harlov, D. E., Förster, H. J., Kotkova, J., & Plasienka, D. (2003). Titanite reaction textures around ilmenite in amphibolite-facies rocks; fluid induced? Journal of the Czech Geological Society = Casopis Ceske Geologicke Spolecnosti, 48(1-2), 57-58.
- Harlov, D. E. (2002). Review of petrographic and mineralogical evidence for fluid induced dehydration of the mafic lower crust; could there be a relationship between granitoids and granulite facies xenoliths in the variscan belt? Zeitschrift für Geologische Wissenschaften, 30(1-2), 13-36.
- Harlov, D. E., & Förster, H. (2002). High-grade fluid metasomatism on both a local and a regional scale; the Seward Peninsula, Alaska, and the Val Strona di Omegna, Ivrea-Verbano zone, northern Italy; part II, phosphate mineral chemistry. Journal of Petrology, 43(5), 801-824. doi:<http://dx.doi.org/10.1093/petrology/43.5.801>
- Harlov, D. E., Wirth, R., & Förster, H. (2005). An experimental study of dissolution-reprecipitation in fluorapatite; fluid infiltration and the formation of monazite. Contributions to Mineralogy and Petrology, 150(3), 268-286. doi:<http://dx.doi.org/10.1007/s00410-005-0017-8>
- Harlov, D., Tropper, P., Seifert, W., Nijland, T., Förster, H., & Baldwin, J. A. (2006). Formation of al-rich titanite (CaTiSiO_4 (sub 4) O-CaAlSiO_4 (sub 4) OH) reaction rims on ilmenite in metamorphic rocks as a function of $f\text{H}_2\text{O}$ and $f\text{O}_2$ (sub 2). Lithos (Oslo), 88(1-4), 72-84. doi:<http://dx.doi.org/10.1016/j.lithos.2005.08.005>
- Hingerl, F., Klötzli, U., Steuber, C., & Kleinschrodt, R. (2008). New results from the mafic complex in the Finero area. International Geological Congress, Abstracts = Congres Geologique International, Resumes, 33, Abstract 1344964.
- Hiraga, T., Anderson, I. M., & Kohlstedt, D. L. (2002). Grain boundary chemistry in mantle rocks. EOS, Transactions, American Geophysical Union, 83(47), 1458.
- Kenkmann, T., & Dresen, G. (2002). Dislocation microstructure and phase distribution in a lower crustal shear zone; an example from the Ivrea-zone, Italy. Geologische Rundschau = International Journal of Earth Sciences (1999), 91(3), 445-458. doi:<http://dx.doi.org/10.1007/s00531-001-0236-9>
- Kissling, E., Bousquet, R., Ford, M., & Schmid, S. (2012). Formation of the arc of the western Alps and Alps-Apennines transition in the light of new geophysical data on the lithospheric architecture around the Ligurian knot. Geophysical Research Abstracts, 14, Abstract EGU2012-8432.
- Kissling, E., Schmid, S. M., Lippitsch, R., Ansorge, J., & Fügenschuh, B. (2006). Lithosphere structure and tectonic evolution of the alpine arc; new evidence from high-resolution teleseismic tomography. Memoirs of the Geological Society of London, 32, 129-145.
- Kleinschrodt, R. (2004). A natural record of lower crustal seismic events. Bochumer Geowissenschaftliche Arbeiten, 3, 172-173.
- Kleinschrodt, R., & Duyster, J. P. (2002). HT-deformation of garnet; an EBSD study on granulites from Sri Lanka, India and the Ivrea zone. Journal of Structural Geology, 24(11), 1829-1844.
- Klötzli, U. S., Sinigoi, S., Quick, J. E., Demarchi, G., Tassinari, C. C. G., Sato, K., & Günes, Z. (2014). Duration of igneous activity in the Sesia magmatic system and implications for high temperature metamorphism in the Ivrea-Verbano deep crust. Lithos (Oslo), 206-207, 19-33. doi:<http://dx.doi.org/10.1016/j.lithos.2014.07.020>
- Kovaleva, E., Klötzli, U., & Habler, G. (2016). Planar deformation bands in zircon; a new evidence of seismicity. International Geological Congress, Abstracts = Congres Geologique International, Resumes, 35, Abstract 111.
- Kovaleva, E., & Klötzli, U. (2017). NanoSIMS study of seismically deformed zircon; evidence of Y, Yb, Ce, and P redistribution and resetting of radiogenic Pb. American Mineralogist, 102(6), 1311-1327. doi:<http://dx.doi.org/10.2138/am-2017-5975>
- Kovaleva, E., Klötzli, U., & Habler, G. (2016). On the geometric relationship between deformation microstructures in zircon and the kinematic framework of the shear zone. Lithos (Oslo), 262, 192-212. doi:<http://dx.doi.org/10.1016/j.lithos.2016.07.001>
- Kovaleva, E., Klötzli, U., Habler, G., Huet, B., Guan, Y., & Rhede, D. (2017). The effect of crystal plastic deformation on isotope and trace element distribution in zircon; combined BSE, CL, EBSD, FEG-EMPA and NanoSIMS study. Chemical Geology, 450, 183-198. doi:<http://dx.doi.org/10.1016/j.chemgeo.2016.12.030>
- Kunz, B. E., Johnson, T. E., White, R. W., & Redler, C. (2014). Partial melting of metabasic rocks in val strona di omegna, Ivrea zone, northern Italy. Lithos (Oslo), 190-191, 1-12. doi:<http://dx.doi.org/10.1016/j.lithos.2013.11.015>
- Kunz, B. E., Manzotti, P., von Niederhäusern, B., Regis, D., Burn, M., Giuntoli, F., & Engi, M. (2015). Tracing Permian metamorphism in fragments of continental crust through rifting, subduction and orogenesis. V.M. Goldschmidt Conference - Program and Abstracts, 25, 1716.

- Langone, A., Renna, M. R., Tiepolo, M., Zanetti, A., Mazzucchelli, M., & Giovanardi, T. (2013). New insights into the evolution of the Finero mafic complex. *Mineralogical Magazine*, 77(5), 1545. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.12>
- Lardeaux, J. M., Schwartz, S., Tricart, P., Paul, A., Guillot, S., Bethoux, N., & Masson, F. (2006). A crustal-scale cross-section of the south-western Alps combining geophysical and geological imagery. *Terra Nova*, 18(6), 412-422. doi:<http://dx.doi.org/10.1111/j.1365-3121.2006.00706.x>
- Lenaz, D., Piani, E., Zaccarini, F., Garuti, G., Princivalle, F., & Raith, J. G. (2005). Spinel geothermometry in the metasomatic chromitites of the Finero mantle massif, Ivrea zone (western Italian Alps). *Mitteilungen Der Österreichischen Mineralogischen Gesellschaft*, 151, 80.
- Lenaz, D., & Princivalle, F. (2011). First occurrence of titanio-magnetites from the websterite dykes within Balmuccia peridotite (Ivrea-Verbano zone); crystal chemistry and structural refinement. *Periodico Di Mineralogia*, 80(1), 19-26. doi:<http://dx.doi.org/10.2451/2011PM0002>
- Locmelis, M., Fiorentini, M. L., Rushmer, T., Adam, J., Zaccarini, F., Garuti, G., . . . Turner, S. (2013). Mass transfer of fluids and metals in the deep earth. *Mineralogical Magazine*, 77(5), 1636. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.12>
- Locmelis, M., Fiorentini, M. L., Rushmer, T., Arevalo, R., Jr., Adam, J., & Denyszyn, S. W. (2016). Sulfur and metal fertilization of the lower continental crust. *Lithos (Oslo)*, 244, 74-93. doi:<http://dx.doi.org/10.1016/j.lithos.2015.11.028>
- Luvizotto, G. L., & Zack, T. (2009). Nb and zr behavior in rutile during high-grade metamorphism and retrogression; an example from the Ivrea-Verbano zone. *Chemical Geology*, 261(3-4), 303-317. doi:<http://dx.doi.org/10.1016/j.chemgeo.2008.07.023>
- Luvizotto, G. L., & Zack, T. (2007). Geochemical trends in rutile forming reactions. *Geochimica Et Cosmochimica Acta*, 71(15), 1.
- Matsumoto, T., Morishita, T., Matsuda, J., Fujioka, T., Takebe, M., Yamamoto, K., & Arai, S. (2005). Noble gases in the Finero phlogopite-peridotites, western Italian Alps. *Earth and Planetary Science Letters*, 238(1-2), 130-145. doi:<http://dx.doi.org/10.1016/j.epsl.2005.07.005>
- Matysiak, A. K., & Trepmann, C. A. (2012). Crystal-plastic deformation and recrystallization of peridotite controlled by the seismic cycle. *Tectonophysics*, 530-531, 111-127. doi:<http://dx.doi.org/10.1016/j.tecto.2011.11.029>
- Matysiak, A., & Trepmann, C. (2014). Cycles of localized deformation and recrystallization in peridotites from the Finero complex (Ivrea zone). *Schriftenreihe Der Deutschen Gesellschaft für Geowissenschaften*, 85, 319.
- Mazzucchelli, M., Zanetti, A., Rivalenti, G., Vannucci, R., Correia, C. T., & Tassinari, C. C. G. (2010). Age and geochemistry of mantle peridotites and diorite dykes from the Baldissaro body; insights into the Paleozoic-Mesozoic evolution of the southern Alps. *Lithos (Oslo)*, 119(3-4), 485-500. doi:<http://dx.doi.org/10.1016/j.lithos.2010.08.002>
- Mulch, A., & Cosca, M. A. (2004). Recrystallization or cooling ages; in situ UV-laser (super 40) Ar/ (super 39) Ar geochronology of muscovite in mylonitic rocks. *Journal of the Geological Society of London*, 161(4), 573- 582. doi:<http://dx.doi.org/10.1144/0016-764903-110>
- Mulch, A., Rosenau, M., Dörr, W., & Handy, M. R. (2002). The age and structure of dikes along the tectonic contact of the Ivrea-Verbano and strona-ceneri zones (southern Alps, northern Italy, Switzerland). *Schweizerische Mineralogische und Petrographische Mitteilungen = Bulletin Suisse De Mineralogie Et Petrographie*, 82(1), 55-76.
- Oberti, R., Boiocchi, M., Hawthorne, F. C., Ball, N. A., Camara, F., Pagano, R., & Pagano, A. (2016). Ferro-ferri- hornblende from the Traversella mine (Ivrea, Italy); occurrence, mineral description and crystal chemistry. *Mineralogical Magazine*, 80(7), 1233-1242. doi:<http://dx.doi.org/10.1180/minmag.2016.080.060>
- Pape, J., Mezger, K., & Robyr, M. (2016). A systematic evaluation of the Zr-in-rutile thermometer in ultrahigh temperature (UHT) rocks. *Contributions to Mineralogy and Petrology*, 171(5), Article 44. doi:<http://dx.doi.org/10.1007/s00410-016-1254-8>
- Paul, A., Cattaneo, M., Thouvenot, F., Spallarossa, D., Bethoux, N., & Frechet, J. (2001). A three-dimensional crustal velocity model of the southwestern Alps from local earthquake tomography. *Journal of Geophysical Research*, 106, 19-19,389. doi:<http://dx.doi.org/10.1029/2001JB000388>
- Peressini, G., Quick, J. E., Sinigoi, S., Hofmann, A. W., & Fanning, M. (2007). Duration of a large mafic intrusion and heat transfer in the lower crust; a SHRIMP U-Pb zircon study in the Ivrea-Verbano zone (western Alps, Italy). *Journal of Petrology*, 48(6), 1185-1218. doi:<http://dx.doi.org/10.1093/petrology/egm014>
- Peressini, G., Mazzucchelli, M., Rivalenti, G., & Hofmann, A. (2004). Triassic U-Pb SHRIMP ages on magmatic zircons from the external gabbro unit of the Finero mafic complex, Ivrea zone, western Italian Alps.
- International Geological Congress, Abstracts = Congres Geologique International, Resumes, 32, Part 2, 1066.

- Pinarelli, L., & Boriani, A. (2007). Tracing metamorphism, magmatism and tectonics in the southern Alps (Italy); constraints from Rb-Sr and Pb-Pb geochronology, and isotope geochemistry. *Periodico Di Mineralogia*, 76(2), 5-24. doi:<http://dx.doi.org/10.2451/2007PM0006>
- Pinarelli, L., Del Moro, A., Boriani, A., & Caironi, V. (2002). Sr, Nd isotope evidence for an enriched mantle component in the origins of the Hercynian gabbro-granite series of the "Serie dei Laghi" (southern Alps, northwestern Italy). *European Journal of Mineralogy*, 14(2), 403-415. doi:<http://dx.doi.org/10.1127/0935-1221/2002/0014-0403>
- Pittarello, L., Habler, G., Abart, R., & Rhede, D. (2015). Garnet growth in frictional melts of the Ivrea zone (Italy). *Italian Journal of Geosciences*, 134(1), 149-161. doi:<http://dx.doi.org/10.3301/IJG.2014.53>
- Pittarello, L., Pennacchioni, G., & Di Toro, G. (2012). Amphibolite facies pseudotachylites in Premosello metagabbro and felsic mylonites (Ivrea zone, Italy). *Tectonophysics*, 580, 43-57. doi:<http://dx.doi.org/10.1016/j.tecto.2012.08.001>
- Qiu, L., Rudnick, R. L., McDonough, W. F., & Bea, F. (2011). The behavior of lithium in amphibolite- to granulite-facies rocks of the Ivrea-Verbano zone, NW Italy. *Chemical Geology*, 289(1-2), 76-85. doi:<http://dx.doi.org/10.1016/j.chemgeo.2011.07.014>
- Quick, J. E., Sinigoi, S., Peressini, G., Demarchi, G., Wooden, J. L., & Sbisa, A. (2009). Magmatic plumbing of a large permian caldera exposed to a depth of 25 km. *Geology (Boulder)*, 37(7), 603-606. doi:<http://dx.doi.org/10.1130/G30003A.1>
- Quick, J. E., Sinigoi, S., Sbisa, A., Demarchi, G., & Richards, I. (2011). Direct observations on the interaction between lower- and upper-crustal magmatic systems in a virtually complete section through 25 km of pre-Alpine crust in the Sesia valley, northwest Italy. American Geophysical Union Fall Meeting, 2011, Abstract V11G-01.
- Quick, J. E., Sinigoi, S., Snoke, A. W., Kalakay, T. J., Mayer, A., & Peressini, G. (2003). Geologic map of the southern Ivrea-Verbano zone, northwestern Italy U. S. Geological Survey, Reston, VA.
- Raffone, N., Le Fevre, B., Zanetti, A., Ottolini, L., Vannucci, R., Rivalenti, G., & Mazzucchelli, M. (2005). Slab- related mantle-metasomatism; a SIMS investigation of LLE composition of Finero phlogopite-peridotite (W Alps). *Ophioliti*, 30(2), 209-210.
- Redler, C., Johnson, T. E., White, R. W., & Kunz, B. E. (2012). Phase equilibrium constraints on a deep crustal metamorphic field gradient; metapelitic rocks from the Ivrea zone (NW Italy). *Journal of Metamorphic Geology*, 30(3), 235-254. doi:<http://dx.doi.org/10.1111/j.1525-1314.2011.00965.x>
- Redler, C. (2013). Partial melting and melt loss; migmatites from Val Strona di Omegna (Ivrea zone, NW Italy). *Mineralogical Magazine*, 77(5), 2036. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.18>
- Redler, C., White, R. W., & Johnson, T. E. (2013). Migmatites in the Ivrea zone (NW Italy); constraints on partial melting and melt loss in metasedimentary rocks from val strona di omegna. *Lithos (Oslo)*, 175-176, 40-53. doi:<http://dx.doi.org/10.1016/j.lithos.2013.04.019>
- Reichelt-Emden-Steuber, C., & Kleinschrodt, R. (2004). The lower crustal profile of the val cannobino (Ivrea zone); magmatic thickening and tectonic telescoping. *Terra Nostra (Bonn)*, 2004-1, 70.
- Rosenberg, C. L., & Kissling, E. (2013). Three-dimensional insight into central-alpine collision; lower-plate or upper-plate indentation? *Geology (Boulder)*, 41(12), 1219-1222. doi:<http://dx.doi.org/10.1130/G34584.1>
- Rutter, E., Brodie, K., James, T., & Burlini, L. (2007). Large-scale folding in the upper part of the Ivrea-Verbano zone, NW Italy. *Journal of Structural Geology*, 29(1), 1-17. doi:<http://dx.doi.org/10.1016/j.jsg.2006.08.013>
- Sacchi, R., Quagliolo, P., & Strona, A. (2006). Il settore sud-occidentale della zona Strona-Ceneri nel Biellese e Canavese. *Bollettino - Museo Regionale Di Scienze Naturali*, 24(1), 5-24.
- Sawaguchi, T., & Ishii, K. (2003). Three-dimensional numerical modeling of lattice- and shape-preferred orientation of orthopyroxene porphyroclasts in peridotites. *Journal of Structural Geology*, 25(9), 1425-1444. doi:[http://dx.doi.org/10.1016/S0191-8141\(02\)00199-2](http://dx.doi.org/10.1016/S0191-8141(02)00199-2)
- Scafidi, D., Solarino, S., Eva, C., & Marson, I. (2006). Structure and properties of the Ivrea body and of the Alps-apennines system as revealed by local earthquake tomography. *Bollettino Di Geofisica Teorica Ed Applicata*, 47(3), 497-514.
- Schaltegger, U., & Brack, P. (2007). Crustal-scale magmatic systems during intracontinental strike-slip tectonics; U, Pb and Hf isotopic constraints from permian magmatic rocks of the southern Alps. *Geologische Rundschau = International Journal of Earth Sciences* (1999), 96(6), 1131-1151. doi:<http://dx.doi.org/10.1007/s00531-006-0165-8>
- Schaltegger, U., Ulianov, A., Müntener, O., Ovtcharova, M., Peytcheva, I., Vonlanthen, P., . . . Girlanda, F. (2015). Megacrystic zircon with planar fractures in miaskite-type nepheline pegmatites formed at high pressures in the lower crust (Ivrea zone, southern Alps, Switzerland). *American Mineralogist*, 100(1), 83-94. doi:<http://dx.doi.org/10.2138/am-2015-4773>
- Selverstone, J., & Sharp, Z. D. (2011). A chlorine isotope view of mantle metasomatism via slab fluids/ melts. *Mineralogical Magazine*, 75(3), 1836.

- Selverstone, J., & Sharp, Z. D. (2011). Chlorine isotope evidence for multicomponent mantle metasomatism in the Ivrea zone. *Earth and Planetary Science Letters*, 310(3-4), 429-440. doi:<http://dx.doi.org/10.1016/j.epsl.2011.08.034>
- Siegesmund, S., Layer, P., Dunkl, I., Vollbrecht, A., Steenken, A., Wemmer, K., & Ahrendt, H. (2008). Exhumation and deformation history of the lower crustal section of the valstrona di omegna in the Ivrea zone, southern Alps. *Geological Society Special Publications*, 298, 45-68. doi:<http://dx.doi.org/10.1144/SP298.3>
- Sinigoi, S., Quick, J. E., Demarchi, G., & Klötzli, U. (2011). The role of crustal fertility in the generation of large silicic magmatic systems triggered by intrusion of mantle magma in the deep crust. *Contributions to Mineralogy and Petrology*, 162(4), 691-707. doi:<http://dx.doi.org/10.1007/s00410-011-0619-2>
- Sinigoi, S. (2007). The Ivrea-Verbano magmatic system. *Acta Vulcanologica*, 19(1-2), 65-66.
- Sinigoi, S., Quick, J. E., Demarchi, G., & Klötzli, U. S. (2016). Production of hybrid granitic magma at the advancing front of basaltic underplating; inferences from the Sesia magmatic system (southwestern Alps, Italy). *Lithos (Oslo)*, 252-253, 109-122. doi:<http://dx.doi.org/10.1016/j.lithos.2016.02.018>
- Sinigoi, S., Quick, J. E., Demarchi, G., & Peressini, G. (2010). The Sesia magmatic system. *Journal of the Virtual Explorer*, 36, 33. doi:<http://dx.doi.org/10.3809/jvirtex.2010.00218>
- Smye, A. J., & Stockli, D. F. (2014). Rutile U-Pb age depth profiling; a continuous record of lithospheric thermal evolution. *Earth and Planetary Science Letters*, 408, 171-182. doi:<http://dx.doi.org/10.1016/j.epsl.2014.10.013>
- Smye, A., Stockli, D. F., & Zack, T. (2014). Tracing lithospheric thermal evolution with high field strength element speedometry in rutile. *American Geophysical Union Fall Meeting*, 2014, Abstract V54B-08, 2.
- Snoke, A. W., & Kalakay, T. J. (2005). A structural transect across the Ivrea-Verbano zone near Varallo Sesia, northern Italy. *Abstracts with Programs - Geological Society of America*, 37(4), 53.
- Solano, J. M. S., Jackson, M. D., Sparks, R. S. J., & Blundy, J. (2014). Evolution of major and trace element composition during melt migration through crystalline mush; implications for chemical differentiation in the crust. *American Journal of Science*, 314(5), 895-939. doi:<http://dx.doi.org/10.2475/05.2014.01>
- Sossi, P. A., O'Neill, H. S., C., & Beltrando, M. (2013). Iron isotope geochemistry of the Balmuccia peridotite massif and the composition of the upper mantle. *Mineralogical Magazine*, 77(5), 2240. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.19>
- Souquiere, F., & Fabbri, O. (2010). Pseudotachylytes in the Balmuccia peridotite (Ivrea zone) as markers of the exhumation of the Southern Alpine continental crust. *Terra Nova*, 22(1), 70-77. doi:<http://dx.doi.org/10.1111/j.1365-3121.2009.00918.x>
- Souquiere, F., Monie, P., Fabbri, O., & Chauvet, A. (2011). Polyphase seismic faulting in the Ivrea zone (Italian Alps) revealed by (super 40) Ar/ (super 39) Ar dating of pseudotachylytes. *Terra Nova*, 23(3), 162-170. doi:<http://dx.doi.org/10.1111/j.1365-3121.2011.00994.x>
- Spalla, M. I., & Zulbati, F. (2004). Structural and petrographic map of the southern Sesia-Lanzo zone; Monte Soglio-Rocca Canavese, western Alps, Italy. *Memorie Di Scienze Geologiche*, 55, 119-127, 1 sheet.
- Stähle, V., Frenzel, G., Hess, J. C., Saupe, F., Schmidt, S. T., & Schneider, W. (2001). Permian metabasalt and triassic alkaline dykes in the northern Ivrea zone; clues to the post-variscan geodynamic evolution of the southern Alps. *Schweizerische Mineralogische Und Petrographische Mitteilungen = Bulletin Suisse De Mineralogie Et Petrographie*, 81(1), 1-21.
- Stepanov, A. S., & Hermann, J. (2013). Biotite is an important host for Nb in the lower crust. *Mineralogical Magazine*, 77(5), 2262. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.19>
- Ueda, T., Obata, M., Ozawa, K., & Di Toro, G. (2009). Ductile and brittle deformation in peridotite recorded in ultramafic pseudotachylytes and associated fault rocks (Balmuccia, Italian Alps). *EOS, Transactions, American Geophysical Union*, 90(52), Abstract V51B-1674.
- Van, d. L., Schüngel, M., Kleinschrodt, R., Münker, C., Kirchenbaur, M., & Fonseca, R. O. C. (2013). Combined Lu-Hf and Sm-Nd garnet-geochronology of lower crustal rocks from Val Strona, Ivrea zone, Italy. *Mineralogical Magazine*, 77(5), 2390. doi:<http://dx.doi.org/10.1180/minmag.2013.077.5.22>
- Vanini, F. (2002). Vanadinite da una cava presso croveo (Baceno, Ossola). *Rivista Mineralogica Italiana*, 26(3), 179.
- Vignola, P., Diella, V., Oppizzi, P., Tiepolo, M., & Weiss, S. (2008). Phosphate assemblages from the Brissago granitic pegmatite, western southern Alps, Switzerland. *The Canadian Mineralogist*, 46(3), 635-650. doi:<http://dx.doi.org/10.3749/canmin.46.3.535>
- Wang, Z., & Becker, H. (2015). Fractionation of highly siderophile and chalcogen elements during magma transport in the mantle; constraints from pyroxenites of the balmuccia peridotite massif. *Geochimica Et Cosmochimica Acta*, 159, 244-263. doi:<http://dx.doi.org/10.1016/j.gca.2015.03.036>
- Wang, Z., Becker, H., & Gawronski, T. (2013). Partial re-equilibration of highly siderophile elements and the chalcogens in the mantle; a case study on the baldissero and balmuccia peridotite massifs (Ivrea zone,

- Italian Alps). *Geochimica Et Cosmochimica Acta*, 108, 21-44. doi:<http://dx.doi.org/10.1016/j.gca.2013.01.021>
- Wawrzenitz, N. H., Romer, R. L., Grasemann, B., & Rhede, D. (2012). Two stages of fluid-rock interaction in UHP marbles (Dabie Shan, China); grain-scale processes and map-scale metasomatism. American Geophysical Union Fall Meeting, 2012, Abstract V43A-2817. Retrieved from <https://search.proquest.com/docview/1549620660?accountid=14682>
 - Weyer, S., Münker, C., & Mezger, K. (2003). Nb/Ta, Zr/Hf and REE in the depleted mantle; implications for the differentiation history of the crust-mantle system. *Earth and Planetary Science Letters*, 205(3-4), 309-324. doi:[http://dx.doi.org/10.1016/S0012-821X\(02\)01059-2](http://dx.doi.org/10.1016/S0012-821X(02)01059-2)
 - Wolff, R., Dunkl, I., Kiesselbach, G., Wemmer, K., & Siegesmund, S. (2012). Thermochronological constraints on the multiphase exhumation history of the Ivrea-Verbano zone of the southern Alps. *Tectonophysics*, 579, 104-117. doi:<http://dx.doi.org/10.1016/j.tecto.2012.03.019>
 - Wu, C. M., & Zhao, G. C. (2006). The applicability of the GRIPS geobarometry in metapelitic assemblages. *Journal of Metamorphic Geology*, 24(4), 297-307. doi:<http://dx.doi.org/10.1111/j.1525-1314.2006.00638.x>
 - Zaccarini, F., Meisel, T., Thalhammer, O. A. R., & Garuti, G. (2005). The Finero chromitites (Ivrea zone, Italy); evidence for metasomatism in a mantle plume.
 - Zack, T., Luvizotto, G. L., Stockli, D. F., & Barth, M. (2008). Texturally controlled U/Pb dating of rutile from the Ivrea zone. *Geochimica Et Cosmochimica Acta*, 72(12), 1. Retrieved from <https://search.proquest.com/docview/50415544?accountid=14682>
 - Zack, T., Stockli, D. F., Luvizotto, G. L., Barth, M. G., Belousova, E., Wolfe, M. R., & Hinton, R. W. (2011). In situ U/Pb rutile dating by LA-ICP-MS; (super 208) pb correction and prospects for geological applications. *Contributions to Mineralogy and Petrology*, 162(3), 515-530. doi:<http://dx.doi.org/10.1007/s00410-011-0609-4>
 - Zaffarana, C. B., Poma, S., & Patino Douce, A. (2009). Thermal effect produced by mafic-ultramafic magma intrusion; an example from Sierra de San Luis, central Argentina. *Abstracts with Programs - Geological Society of America*, 41(7), 147.
 - Zanetti, A., Mazzucchelli, M., Giovanardi, T., Tiepolo, M., & Vannucci, R. (2011). Mantle migration of K-LILE-enriched melts/fluids in supra-subduction settings; evidence from the Finero complex (Ivrea-Verbano zone, southern Alps). American Geophysical Union Fall Meeting, 2011, Abstract V41D-2538. Retrieved from <https://search.proquest.com/docview/1696872931?accountid=14682>
 - Zanetti, A., Giovanardi, T., Langone, A., Tiepolo, M., Wu, F., Dallai, L., & Mazzucchelli, M. (2016). Origin and age of zircon-bearing chromitite layers from the Finero phlogopite peridotite (Ivrea-Verbano zone, western Alps) and geodynamic consequences. *Lithos (Oslo)*, 262, 58-74. doi:<http://dx.doi.org/10.1016/j.lithos.2016.06.015>
 - Zanetti, A., Mazzucchelli, M., Sinigoi, S., Giovanardi, T., Peressini, G., & Fanning, M. (2013). SHRIMP U-Pb zircon triassic intrusion age of the Finero mafic complex (Ivrea-Verbano zone, western Alps) and its geodynamic implications. *Journal of Petrology*, 54(11), 2235-2265. doi:<http://dx.doi.org/10.1093/petrology/egt046>
 - Zappone, A. S., & Benson, P. M. (2013). Effect of phase transitions on seismic properties of metapelites; a new high-temperature laboratory calibration. *Geology (Boulder)*, 41(4), 463-466. doi:<http://dx.doi.org/10.1130/G33713.1>
 - Zhong, X., Kunze, K., & Zappone, A. S. (2013). Can minor hydrous minerals play a role in low crust seismic anisotropy; an experimental and numerical investigation to Finero peridotites. American Geophysical Union Fall Meeting, 2013, Abstract MR43A-2391. Retrieved from <https://search.proquest.com/docview/1692746352?accountid=14682>
 - Zhong, X., Frehner, M., Kunze, K., & Zappone, A. (2014). A novel EBSD-based finite-element wave propagation model for investigating seismic anisotropy; application to Finero peridotite, Ivrea-Verbano zone, northern Italy. *Geophysical Research Letters*, 41(20), 7105-7114. doi:<http://dx.doi.org/10.1002/2014GL060490>
 - Zucali, M., Barberini, V., Voltolini, M., Ouladdiaf, B., Chateigner, D., Mancini, L., & Lutterotti, L. (2014). Quantitative 3D microstructural analysis of naturally deformed amphibolite from the southern Alps (Italy); microstructures, CPO and seismic anisotropy from a fossil extensional margin. *Special Publication - Geological Society of London*, 409(1), 201-222, 22. doi:<http://dx.doi.org/10.1144/SP409.5>